



1

Galton

1875 Francis

" 973"

>>>

"

2014 11

"

2

twin study

nature versus nurture

monozygotic MZ

dizygotic DZ

50%

MZ DZ

1

candidate gene association study CGAS

single

nucleotide polymorphism SNP SNP

DNA

90% 500 1000

2014-2015

72

The figure is a scatter plot with 'variable number of tandem repeats VNTR' on the x-axis (ranging from 0 to 300) and 'haplotype' on the y-axis (ranging from 0 to 10%). Blue diamond-shaped data points show a positive correlation with the red line. A horizontal dashed line is drawn across the plot at approximately 10% on the y-axis.

—

SNP

2                    3                    5

24                    rs6658108            rs732050


rs1487275            rs8076005            rs2192371

19%

```

graph TD
    A[genome-wide association study] --> B[SNP]
    A --> C[copy number variation]
    B --> D[CGAS]
    B --> E[GWAS]
    C --> F[GWAS]
  
```


>>>



2014–2015



fMRI和APOE ( Bookheimer et al., 2001 ) /  
COMT ( Egan et al., 2001 ) /5-HTTLPR  
( Hariri et al., 2002 )：  
第一次基于MRI的神经影像遗传学



1 10

case-control study

family study

animal study

knockout

3

2001

Chinese National Twin Registry CNTR


TRiSC

The Beijing Twin Study BeTwiSt

20

Twin Registry in Southwestern China

21



|||||

generalist genes

22

FRA

FIE

CFE

23

24

25

Val66Met

26 27

SLEs

Val

MetMet

28

50%

30%

29

2

SNP rs42352

RAPM

30

rs7435958

31

CT/TT

32-35

35

rs4334545

C

rs4334545

CC

&gt;&gt;&gt;

表2 国内认知能力的遗传基础研究的主要进展

30	169 SNP	RAPM	SNP rs42352 rs42352 TT RAPM
31	49 SNP	WAIS-R	rs7435958 GG rs7435958 GG
32	Val158Met rs4680 COMT		Met Val 2-back
35	5 SNP		2-back      rs4334545      C rs6090453      G rs4334545 CT/TT CC
33	Val158Met 1403C/ T - 48A/G TaqIA TaqIB Ser9Gly - C521T - G809A VNTR		TaqIA      TaqIB Ser9Gly TaqIA
34	rs6265 Val 166Met 5-		Val BDNF Met
42	7 DDC 5-		SNP      rs3887825      rs7786398 rs10499695      rs6969081 rs7786398      rs10499695      rs6969081 SNP G rs10499695      rs6969081
			st      798 Val BDNF

37	- 1021C/T 19bp Ins/Del 30bp VNTR rs1799836		- 19bp Ins/Del VNTR
40	VNTR A	5-	4r/4r 3r/3r 3r/4r " "
39			AAGG/AAGG SSRT Stop-signal reaction time CCAC/AAGG CCAC/CCAC AAGG/AAGG
41	30 SNP 5- 2A		rs1923888 rs1745837 rs9567739 rs3742279 rs655888 rs655854 and rs2296972
38	13	5-	T267C T267C TT CT TT

36-42

3

3

18%

43

5- HTTLPR

44 45

79

表3 国内情绪及社会行为的遗传基础研究的主要进展

3	127 SNP 2 VNTR 5		8 5- 12 SNP HTR6 rs6658108 HTR5A rs732050 TPH2 rs1487275 s8076005 rs2192371 19%
45	5- 5 HTTLPR 5		5-HTTLPR l/l s/s
44	5- 5 HTTLPR 5		5-HTTLPR l/l s/s HA 5-HTTLPR HA
57	5- 5 HTTLPR 5	5-	l/l 5- s/s
46	rs53576		rs53576 A rs53576 rs53576


47	rs53576		A/A A/A	G/G G/G
48	rs53576		A/A G/G	G A/A
43	96 SNP 2			18%
58	5-HTTLPR 5	5	5-HTTLPR 1 LAT	IGT S IGT 40
59	Val158Met rs4680			

>>>

50	C- 1019G rs6295 5- 1A		CG/GG	CC
49	- 1021C/T 19bp Ins/Del		- 1021C/T CC DBH CT/TT 2.3% 1.4%	
61	10	IRI	O rs2254298 rs2268491 rs13316193 rs4686302 IRI	
62	7 SNP COMT		rs2020917 and rs737865 SNP rs5993883 SNP	


### 3.4.1 注意缺陷多动障碍


attention deficit hyperactivity disorder ADHD




### 3.4.2 重度抑郁症

major depression disorder MDD





#### 3.4.4 双相情感障碍





|||||

4

22

90

138

139

W. S. Kremen

140 141

J. L. Stein      GWAS

142

80%

GWAS

143-145

GWAS

ENIGMA   CHARGE   IMAGEN

146

5

85

>>>

1

5 10

2

GWAS

GWAS

3

GWAS

4

-

147

5

5 10

2014–2015

## 部分生物学专业名词解释

genetic material	RNA DNA		
heritability			
gene polymorphism		2	
allele			
genotype			
haplotype			
copy number variation CNV		1 kb	
variable number of tandem repeats VNTR	10    1000	DNA	10
linkage			
phenotype			
gene frequency			
susceptibility			

## —— 参考文献 ——

- 1 J 2012 44 3  
331- 333
- 2 De Quervain D J, Papassotiropoulos A. Identification of a genetic cluster influencing memory performance and hippocampal activity in humans J . Proc Natl Acad Sci U S A, 2006, 103 11 4270-4274.
- 3 Chen C, Liu C, Chen C, et al. Genetic variations in the serotonergic system and environmental factors contribute to aggressive behavior in Chinese adolescents J . Physiol Behav, 2015, 138 62-68
- 4 Lee S H, Wray N R, Goddard M E, et al. Estimating missing heritability for disease from genome-wide association studies J . The American Journal of Human Genetics, 2011, 88 3 294-305.
- 5 Casamassima F, Hay A C, Benedetti A, et al. L<sup>+</sup> type calcium channels and psychiatric disorders A brief review J .

- American Journal of Medical Genetics Part B Neuropsychiatric Genetics, 2010, 153 8 1373-1390
- 6 Ott J, Kamatani Y, Lathrop M. Family-based designs for genome-wide association studies J. Nature reviews Genetics, 2011, 12 7 465-474
  - 7 Suo C, Touloupou T, Bramon E, et al. Analysis of multiple phenotypes in genome-wide genetic mapping studies J. BMC bioinformatics, 2013, 14 151.
  - 8 Zhang K, Chang S, Cui S, et al. ICSNPathway identify candidate causal SNPs and pathways from genome-wide association study by one analytical framework J. Nucleic acids research, 2011, 39 Web Server issue W437-443
  - 9 Zhang K, Cui S, Chang S, et al. i-GSEA4GWAS a web server for identification of pathways/gene sets associated with traits by applying an improved gene set enrichment analysis to genome-wide association study J. Nucleic acids research, 2010, 38 Web Server issue W90-95
  - 10 Bigos K L, Weinberger D R. Imaging genetics - days of future past J. Neuroimage, 2010, 53 3 804-809
  - 11 Heinz A, Goldman D, Jones D W, et al. Genotype influences in vivo dopamine transporter availability in human striatum J. Neuropsychopharmacol, 2000, 22 2 133-139.
  - 12 Buckhimer S Y, Strijwas MH, Chen MS, et al. Patterns of brain activation in people at risk for Alzheimer's disease J. N Engl J Med, 2000, 343 7 450-456
  - 13 Egan MF, Goldberg TE, Kolachana B S, et al. Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia J. Proc Natl Acad Sci U S A, 2001, 98 12 6917-6922
  - 14 Hariri A R, Mattay V S, Tessitore A, et al. Serotonin transporter genetic variation and the response of the human amygdala J. Science, 2002, 297 5580 400-403
  - 15 Meyer-Lindenberg A, Nichols T, Callicott J H, et al. Impact of complex genetic variation in COMT on human brain function J. Mol Psychiatry, 2006, 11 9 867-877, 797.
  - 16 Potkin S G, Turner J A, Fallon J A, et al. Gene discovery through imaging genetics identification of two novel genes associated with schizophrenia J. Mol Psychiatry, 2009, 14 4 416-428
  - 17 Pezawas L, Meyer-Lindenberg A, Goldman A L, et al. Evidence of bidirectional epistasis between BDNF and SLC6A4 and implications for depression J. Mol Psychiatry, 2008, 13 7 709-716
  - 18 Esslinger C, Walter H, Kirsch P, et al. Neural mechanisms of a genome-wide supported psychosis variant J. Science, 2009, 324 5927 605.
  - 19 Nicodemus K, Law A J, Radulescu E, et al. Biological validation of b s f dhalf cphrenia J. ? / w  
im6 QScien ssl" ssfo ?" l. l. " agh ? gen  
claim

- 27 Martinowich K, Manji H, Lu B. New insights into BDNF function in depression and anxiety. *J . Nat Neurosci*, 2007, 10 9 1089-1093
- 28 Chen J, Li X, McGue M. The interacting effect of the BDNF Val66Met polymorphism and stressful life events on adolescent depression is not an artifact of gene-environment correlation: evidence from a longitudinal twin study. *J . J Child Psychol Psychiatry*, 2013, 54 10 1066-1073.
- 29 Deary I J, Johnson W, Houlihan L M. Genetic foundations of human intelligence. *J . Hum Genet*, 2009, 126 1 215-232
- 30 Zhu B, Chen C, Xue G, et al. The SEMA5A gene is associated with hippocampal volume, and their interaction is associated with performance on Raven's Progressive Matrices. *J . Neuroimage*, 2013, 88C 181-187.
- 31 Zhu B, Chen C, Xue G, et al. The GABRB1 gene is associated with thalamus volume and modulates the association between thalamus volume and intelligence. *J . Neuroimage*, 2014, 102 Pt 2 756-763
- 32 Wang Y, Li J, Chen C, et al. COMT rs4680 Met is not always the 'smart allele': Val allele is associated with better working memory and larger hippocampal volume in healthy Chinese. *J . Genes Brain Behav*, 2013, 12 3 323-329
- 33 Gong P, Zhang H, Chi W, et al. An association study on the polymorphisms of dopaminergic genes with working memory in a healthy Chinese Han population. *J . Cell Mol Neurobiol*, 2012, 32 6 1011-1019
- 34 Gong P, Zheng A, Chen D, et al. Effect of BDNF Val66Met polymorphism on digital working memory and spatial localization in a healthy Chinese Han population. *J . J Mol Neurosci*, 2009, 38 3 250-256
- 35 Li J, Chen C, Chen C, et al. Neuregulin receptor 1 gene NTR1 polymorphism is associated with working memory. *J . PLoS One*, 2011, 6 3 e17365
- 36 Gong P, Xi S, Li S, et al. Effect of Val66Met polymorphism in BDNF on attentional bias in an extroverted Chinese Han population. *J . Acta Neuropathol Exp Wars*, 2013, 73 2 280-288
- 37 Gong P, Xi S, Shen G, et al. The effects of DBH, MAOA, and MAOB on attentional biases for facial expressions. *J . J Mol Neurosci*, 2013, 49 3 606-613
- 38 Gong P, Zheng Z, Chi W, et al. An association study of the genetic polymorphisms in 13 neural plasticity-related genes with semantic and episodic memories. *J . J Mol Neurosci*, 2012, 46 2 352-361.
- 39 Lei X, Chen C, He Q, et al. Haplotype polymorphism in the alpha-2B-adrenergic receptor gene influences response inhibition in a large Chinese sample. *J . Neuropsychopharmacol*, 2012, 37 5 1115-1121.
- 40 Zhang M, Chen X, Way N, et al. The association between infants' self-regulatory behavior and MAOA gene polymorphism. *J . Dev Sci*, 2011, 14 5 1059-1065
- 41 Zhu B, Chen C, Loftus E F, et al. True but not false memories are associated with the HTR2A gene. *J . Neurobiol Learn Mem*, 2013, 106 204-209
- 42 Zhu B, Chen C, Moysis R K, et al. The DOPA decarboxylase DDC gene is associated with alerting attention. *J . Prog Neuropsychopharmacol Biol Psychiatry*, 2013, 43 140-145
- 43 Zhu B, Chen C, Moysis R K, et al. Genetic variations in the dopamine system and facial expression recognition in healthy Chinese college students. *J . Neuropsychobiology*, 2012, 65 2 83-89
- 44 Ma Y, Li B, Wang C, et al. 5-HTTLPR polymorphism modulates neural mechanisms of negative self-reflection. *J . Cereb Cortex*, 2014, 24 9 2421-2429
- 45 Ma Y, Wang C, Li B, et al. Does self-construal predict activity in the social brain network? A genetic moderation effect. *J . Soc Cogn Affect Neurosci*, 2014, 9 9 1360-1367.
- 46 Luo S, Han S. The association between an oxytocin receptor gene polymorphism and cultural orientations. *J . Culture and Brain*, 2014, 2 1 89-107.
- 47 Luo S, Li B, Ma Y, et al. Oxytocin receptor gene and racial ingroup bias in empathy-related brain activity. *J . Neuroimage*, 2015, 110 22-31.
- 48 Luo S, Ma Y, Liu Y, et al. Interaction between oxytocin receptor polymorphism and interdependent culture values on

- human empathy J . Social cognitive and affective neuroscience, 2015, nsv019.
- 49 Gong P, Liu J, Li S, et al. Dopamine beta-hydroxylase gene modulates individuals' empathic ability J . Soc Cogn Affect Neurosci, 2014, 9 9 1341- 1345.
- 50 Gong P, Liu J, Li S, et al. Serotonin receptor gene 5-HT1A modulates alexithymic characteristics and attachment orientation J . Psychoneuroendocrin, 2014, 50 274-279.
- 51 Liu J, Gong P, Zhou X. The association between romantic relationship status and 5-HT1A gene in young adults J . Sci Rep, 2014, 4 7049.
- 52 Li M, Chen J, Li N, et al. A twin study of problematic internet use its heritability and genetic association with effortful control J . Twin Res HumGenet, 2014, 17 4 279-287.
- 53 Chow B W, Ho C S, Wong S W, et al. Generalist genes and cognitive abilities in Chinese twins J . Dev Sci, 2013, 16 2 260-268.
- 54 Hou J, Chen Z, Natsuaki M N, et al. A longitudinal investigation of the associations among parenting, deviant peer affiliation, and externalizing behaviors a monozygotic twin differences design J . Twin Res HumGenet, 2013, 16 3 698-706.
- 55 Chen J, Li X. Genetic and environmental etiologies of adolescent dysfunctional attitudes a twin study J . Twin Res HumGenet, 2014, 17 1 16-22.
- 56 Chen J, Yu J, Li X, et al. Genetic and environmental contributions to anxiety among Chinese children and adolescents - a multi-informant twin study J . J Child Psychol Psychiatry, 2014.
- 57 Ma Y, Li B, Wang C, et al. Allelic variation in 5-HTTLPR and the effects of citalopram on the emotional neural network J . The British Journal of Psychiatry, 2015, 206 5 385-392.
- 58 He Q, Xue G, Chen C, et al. Serotonin transporter gene-linked polymorphic region 5-HTTLPR influences decision making under ambiguity and risk in a large Chinese sample J . Neuropharmacology, 2010, 59 6 518-526.
- 59 He Q, Xue G, Chen C, et al. COMT Val158Met polymorphism interacts with stressful life events and parental warmth to influence decision making J . Sci Rep, 2012, 2 677.
- 60 Lei X, Chen C, He Q, et al. Sex determines which section of the SLC6A4 gene is linked to obsessive-compulsive symptoms in normal Chinese college students J . J Psychiatr Res, 2012, 46 9 1153-1160.
- 61 Wu N, Li Z, Su Y. The association between oxytocin receptor gene polymorphism OXTR and trait empathy J . J Affect Disord, 2012, 138 3 468-472.
- 62 Xia H, Wu N, Su Y. Investigating the genetic basis of theory of mind ToM the role of catechol-O-methyltransferase COMT gene polymorphisms J . PLoS One, 2012, 7 11 e49768.
- 63 Wood A C, Rijssdijk F, Saudino K J, et al. High heritability for a composite index of children's activity level measures J . Behav Genet, 2008, 38 3 266-276.
- 64 Gizer I R, Ficks C, Waldman I D. Candidate gene studies of ADHD a meta-analytic review J . HumGenet, 2009, 126 1 51-90.
- 65 Guan L, Wang B, Chen Y, et al. A high-density single-nucleotide polymorphism screen of 23 candidate genes in attention deficit hyperactivity disorder suggesting multiple susceptibility genes among Chinese Han population J . Mol Psychiatry, 2009, 14 5 546-554.
- 66 Qian Q J, Yang L, Wang Y F, et al. Gene-gene interaction between COMT and MAOA potentially predicts the intelligence of attention-deficit hyperactivity disorder boys in China J . Behav Genet, 2010, 40 3 357-365.
- 67 Li H, Liu L, Tang Y, et al. Sex-specific association of brain-derived neurotrophic factor BDNF Val66Met polymorphism and plasma BDNF with attention-deficit/hyperactivity disorder in a drug-naïve Han Chinese sample J . Psychiatry Res, 2014, 217 3 191-197.
- 68 Liu L, Guan L L, Chen Y, et al. Association analyses of MAOA in Chinese Han subjects with attention-deficit/hyperactivity disorder family-based association test, case-control study, and quantitative traits of impulsivity J .

- Am J Med Genet B Neuropsychiatr Genet, 2011, 156B 6 737-748
- 69 Li Y, Baker- Ericzen M, Ji N, et al. Do SNPs of DRD4 gene predict adult persistence of ADHD in a Chinese sample J . Psychiatry Res, 2013, 205 1-2 143-150
- 70 Yang L, Qian Q, Liu L, et al. Adrenergic neurotransmitter system transporter and receptor genes associated with atomoxetine response in attention-deficit hyperactivity disorder children J . J Neural Transm, 2013, 120 7 1127-1133
- 71 Wu J, Xiao H, Sun H, et al. Role of dopamine receptors in ADHD - a systematic meta-analysis J . Mol Neurobiol, 2012, 45 3 605-620
- 72 Sullivan P F, Neale M C, Kendler K S. Genetic epidemiology of major depression - review and meta-analysis J . The American journal of psychiatry, 2000, 157 10 1552-1562
- 73 Lu B, Gottschalk W. Modulation of hippocampal synaptic transmission and plasticity by neurotrophins J . Prog Brain Res, 2000, 128 231-241.
- 74 Nibuya M, Morinobu S, Duman R S. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments J . J Neurosci, 1995, 15 11 7539-7547.
- 75 Chen J, Li X, McGue M. Interacting effect of BDNF Val66Met polymorphism and stressful life events on adolescent depression J . Genes brain and behavior, 2012
- 76 Yang C, Xu Y, Sun N, et al. The combined effects of the BDNF and GSK3B genes modulate the relationship between negative life events and major depressive disorder J . Brain research, 2010, 1355 1-6
- 77 Li Z, Zhang Y, Wang Z, et al. The role of BDNF, NTRK2 gene and their interaction in development of treatment-resistant depression - data from multicenter, prospective, longitudinal clinic practice J . Journal of psychiatric research, 2013, 47 1 8-14
- 78 Xiao Z, Liu W, Gao K, et al. Interaction between CRHR1 and BDNF genes increases the risk of recurrent major depressive disorder in Chinese population J . PLoS One, 2011, 6 12 e28733
- 79 Zhang K, Xu Q, Xu Y, et al. The combined effects of the 5-HTTLPR and 5-HT1A genes modulates the relationship between negative life events and major depressive disorder in a Chinese population J . J Affect Disord, 2009, 114 1-3 224-231.
- 80 Gao Z, Yuan H, Sun M, et al. The association of serotonin transporter gene polymorphism and geriatric depression - a meta-analysis J . Neurosci Lett, 2014, 578 148-152
- 81 Fang J, Yan W, Jiang G X, et al. Serotonin transporter gene polymorphism in Chinese patients with poststroke depression - a case-control study J . Stroke, 2011, 42 5 1461-1463
- 82 Xu H, Zhang Q, Hou X, et al. The effect of the polymorphisms of 5-HTTLPR on new onset of depression in patients who underwent pacemaker implantation J . Psychiatr Genet, 2014, 24 2 70-74
- 83 Fang Z, Zhu S, Gillihan S J, et al. Serotonin transporter genotype modulates functional connectivity between amygdala and PCC/PCu during mood recovery J . Front Hum Neurosci, 2013, 7 704
- 84 Gao J, Pan Z, Jiao Z, et al. TPH2 gene polymorphisms and major depression - a meta-analysis J . PLoS One, 2012, 7 5 e36721.
- 85 Domschke K, Dannlowski U. Imaging genetics of anxiety disorders J . Neuroimage, 2010, 53 3 822-831.
- 86 Chi S, Teng L, Song J H, et al. Tryptophan hydroxylase 2 gene polymorphisms and poststroke anxiety disorders J . J Affect Disord, 2013, 144 1-2 179-182
- 87 Long H, Liu B, Hou B, et al. The long rather than the short allele of 5-HTTLPR predisposes Han Chinese to anxiety and reduced connectivity between prefrontal cortex and amygdala J . Neurosci Bull, 2013, 29 1 4-15
- 88 Zhang L, Liu L, Li X, et al. Serotonin transporter gene polymorphism 5-HTTLPR influences trait anxiety by modulating the functional connectivity between the amygdala and insula in Han Chinese males J . Hum Brain Mapp, 2015.

- 89 Cadore R J. Evidence for genetic inheritance of primary affective disorder in adoptees J . AmJ Psychiatry, 1978 135 4 463- 466
- 90 Xu J, Liu Y, Wang P, et al. Positive association between the brain- derived neurotrophic factor BDNF gene and bipolar disorder in the Han Chinese population J . AmJ Med Genet B Neuropsychiatr Genet, 2010, 153B 1 275- 279.
- 91 Wang Z, Li Z, Chen J, et al. Association of BDNF gene polymorphism with bipolar disorders in Han Chinese population J . Genes Brain Behav, 2012, 11 5 524- 528
- 92 Wang Z, Zhang C, Huang J, et al. MiRNA- 206 and BDNF genes interacted in bipolar I disorder J . J Affect Disord, 2014, 162 116- 119.
- 93 Jiang H Y, Qiao F, Xu X F, et al. Meta- analysis confirms a functional polymorphism 5-HTTLPR in the serotonin transporter gene conferring risk of bipolar disorder in European populations J . Neurosci Lett, 2013, 549 191- 196
- 94 Chang S H, Gao L, Li Z, et al. BDgene a genetic database for bipolar disorder and its overlap with schizophrenia and major depressive disorder J . Biological psychiatry, 2013, 74 10 727- 733
- 95 Li X, Hu Z, He Y, et al. Association analysis of CNTNAP2 polymorphisms with autism in the Chinese Han population J . Psychiatr Genet, 2010, 20 3 113- 117.
- 96 Zhou X, Xu Y, Wang J, et al. Replication of the association of a MET variant with autism in a Chinese Han population J . PLoS One, 2011, 6 11 e27428
- 97 Liu Y, Hu Z, Xun G, et al. Mutation analysis of the NRXN1 gene in a Chinese autism cohort J . J Psychiatr Res, 2012, 46 5 630- 634
- 98 Guo T, Chen H, Liu B, et al. Methylenetetrahydrofolate reductase polymorphisms C677T and risk of autism in the Chinese Han population J . Genet Test Mol Biomarkers, 2012, 16 8 968- 973
- 99 Bi C, Wu J, Jiang T, et al. Mutations of ANK3 identified by exome sequencing are associated with autism susceptibility J . HumMutat, 2012, 33 12 1635- 1638
- 100 Yang Y, Pan C. Role of metabotropic glutamate receptor 7 in autism spectrum disorders a pilot study J . Life Sci, 2013, 92 2 149- 153
- 101 Tian P. RELN gene polymorphisms and susceptibility to autism in Chinese Han population J . Neurul India, 2012, 60 6 581- 584
- 102 Wang L, Li J, Ruan Y, et al. Sequencing ASMT identifies rare mutations in Chinese Han patients with autism J . PLoS One, 2013, 8 1 e53727.
- 103 Shao S, Xu S, Yang J, et al. A commonly carried genetic variant, rs9616915, in SHANK3 gene is associated with a reduced risk of autism spectrum disorder replication in a Chinese population J . Mol Biol Rep, 2014, 41 3 1591- 1595
- 104 Xu X, Xiong Z, Zhang L, et al. Variations analysis of NLGN3 and NLGN4X gene in Chinese autism patients J . Mol Biol Rep, 2014, 41 6 4133- 4140
- 105 Liang S, Wang X L, Zou M Y, et al. Family- based association study of ZNF533, DOCK4 and IMMP2L gene polymorphisms linked to autism in a northeastern Chinese Han population J . J Zhejiang Univ Sci B, 2014, 15 3 264- 271.
- 106 Lu L, Guo H, Peng Y, et al. Common and rare variants of the THBS1 gene associated with the risk for autism J . Psychiatr Genet, 2014, 24 6 235- 240
- 107 Zheng F, Wang L, Jia M, et al. Evidence for association between Disrupted- in- schizophrenia 1 DISC1 gene polymorphisms and autism in Chinese Han population a family- based association study J . Behav Brain Funct, 2011, 7 1 14
- 108 Li J, Liu J, Zhao L, et al. Association study between genes in Reelin signaling pathway and autism identifies DAB1 as a susceptibility gene in a Chinese Han population J . Progress in Neuro- Psychopharmacology and Biological Psychiatry, 2013, 44 226- 232

- 109 Yang W, Liu J, Zheng F, et al. The evidence for association of ATP2B2 polymorphisms with autism in Chinese Han population J . PloS one, 2013, 8 4 e61021.
- 110 Xu L M, Li J R, Huang Y, et al. AutismKB – an evidence-based knowledgebase of autism genetics J . Nucleic acids research, 2012, 40 Database issue D1016-1022.
- 111 Sullivan P F, Kendler K S, Neale M C. Schizophrenia as a complex trait – evidence from a meta-analysis of twin studies J . Arch Gen Psychiat, 2003, 60 12 1187-1192.
- 112 Li M, Luo X J, Xiao X, et al. Allelic differences between Han Chinese and Europeans for functional variants in ZNF804A and their association with schizophrenia J . Am J Psychiatry, 2011, 168 12 1318-1325.
- 113 Li M, Shi C J, Shi Y Y, et al. ZNF804A and schizophrenia susceptibility in Asian populations J . Am J Med Genet B Neuropsychiatr Genet, 2012, 159B 7 794-802.
- 114 Li M, Zhang H, Luo X J, et al. Meta-analysis indicates that the European GWAS-identified risk SNP rs1344706 within ZNF804A is not associated with schizophrenia in Han Chinese population J . PLoS One, 2013, 8 6 e65780.
- 115 Yang Y, Li W, Yang G, et al. Evaluation of the relationship between the ZNF804A single nucleotide polymorphism rs1344706 A/C variant and schizophrenia subtype in Han Chinese patients J . Int J Psychiatry Med, 2013, 45 3 269-278.
- 116 Zhou D H, Yan Q Z, Yan X M, et al. The study of BDNF Val66Met polymorphism in Chinese schizophrenic patients J . Prog Neuropsychopharmacol Biol Psychiatry, 2010, 34 6 930-933.
- 117 Yi Z, Zhang C, Wu Z, et al. Lack of effect of brain derived neurotrophic factor BDNF Val66Met polymorphism on early onset schizophrenia in Chinese Han population J . Brain Res, 2011, 1417 146-150.
- 118 Lu W, Zhang C, Yi Z, et al. Association between BDNF Val66Met polymorphism and cognitive performance in antipsychotic-naïve patients with schizophrenia J . J Mol Neurosci, 2012, 47 3 505-510.
- 119 Li W, Zhou N, Yu Q, et al. Association of BDNF gene polymorphisms with schizophrenia and clinical symptoms in a Chinese population J . Am J Med Genet B Neuropsychiatr Genet, 2013, 162B 6 538-545.
- 120 Zhai J, Yu Q, Chen M, et al. Association of the brain-derived neurotrophic factor gene G196A rs6265 polymorphisms and the cognitive function and clinical symptoms of schizophrenia J . Int J Clin Exp Pathol, 2013, 6 8 1617-1623.
- 121 Wang Y, Fang Y, Shen Y, et al. Analysis of association between the catechol-O-methyltransferase COMT gene and negative symptoms in chronic schizophrenia J . Psychiatry Res, 2010, 179 2 147-150.
- 122 Li W J, Kou C G, Yu Y, et al. Association of catechol-O-methyltransferase gene polymorphisms with schizophrenia and negative symptoms in a Chinese population J . Am J Med Genet B Neuropsychiatr Genet, 2012, 159B 4 370-375.
- 123 Liu X, Hong X, Chan R C, et al. Association study of polymorphisms in the alpha 7 nicotinic acetylcholine receptor subunit and catechol-O-methyl transferase genes with sensory gating in first-episode schizophrenia J . Psychiatry Res, 2013, 209 3 431-438.
- 124 Hui L, Zhang X, Huang X F, et al. The dopamine b-hydroxylase 19 bp insertion/deletion polymorphism was associated with first-episode but not medicated chronic schizophrenia J . J Psychiatr Res, 2012, 46 6 733-737.
- 125 Hui L, Zhang X, Yu Y Q, et al. Association between DBH 19 bp insertion/deletion polymorphism and cognition in first-episode schizophrenic patients J . Schizophr Res, 2013, 147 2-3 236-240.
- 126 Lai J H, Zhu Y S, Huo Z H, et al. Association study of polymorphisms in the promoter region of DRD4 with schizophrenia, depression, and heroin addiction J . Brain Res, 2010, 1359 227-232.
- 127 Zhang F, Fan H, Xu Y, et al. Converging evidence implicates the dopamine D3 receptor gene in vulnerability to schizophrenia J . Am J Med Genet B Neuropsychiatr Genet, 2011, 156B 5 613-619.
- 128 Xiao L, Shen T, Peng D H, et al. Functional -141C Ins/Del polymorphism in the dopamine D2 receptor gene

- promoter and schizophrenia in a Chinese Han population J . J Int Med Res, 2013, 41 4 1171- 1178
- 129 Pan Y, Yao J, Wang B. Association of dopamine D1 receptor gene polymorphism with schizophrenia a meta-analysis J . Neuropsychiatr Dis Treat, 2014, 10 1133- 1139.
- 130 Wei Y L, Li C X, Li S B, et al. Association study of monoamine oxidase A/B genes and schizophrenia in Han Chinese J . Behav Brain Funct, 2011, 7 42.
- 131 Sun Y, Zhang J, Yuan Y, et al. Study of a possible role of the monoamine oxidase A MAOA gene in paranoid schizophrenia among a Chinese population J . Am J Med Genet B Neuropsychiatr Genet, 2012, 159B 1 104-111.
- 132 Zhang R, Zhang H, Li M, et al. Genetic analysis of common variants in the CMY45 cardiomypathy- associated 5 gene with schizophrenia J . Prog Neuropsychopharmacol Biol Psychiatry, 2013, 46 64-69.
- 133 Gu L Z, Jiang T, Cheng Z H, et al. rs11098403 polymorphism near NDST3 is associated with a reduced risk of schizophrenia in a Han Chinese population J . Neurosci Lett, 2014, 581 42-45.
- 134 Luo X J, Li M, Huang L, et al. Convergent lines of evidence support CAMKK2 as a schizophrenia susceptibility gene J . Mol Psychiatry, 2014, 19 7 774-783.
- 135 Wong E H, So H C, Li M, et al. Common variants on Xq28 conferring risk of schizophrenia in Han Chinese J . Schizophr Bull, 2014, 40 4 777-786.
- 136 Zhang B, Xu Y H, Wei S G, et al. Association study identifying a new susceptibility gene AUTS2 for schizophrenia J . International journal of molecular sciences, 2014, 15 11 19406-19416.
- 137 Ji W, Li T, Pan Y, et al. CNTNAP2 is significantly associated with schizophrenia and major depression in the Han Chinese population J . Psychiatry research, 2013, 207 3 225-228.
- 138 Aimone J B, Li Y, Lee S W, et al. Regulation and function of adult neurogenesis from genes to cognition J . Physiol Rev, 2014, 94 4 991-1026.
- 139 Egan MF, Kojima M, Callicott J H, et al. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function J . Cell, 2003, 112 2 257-269.
- 140 Kremen WS, Prom-Wiemer E, Panizzon M S, et al. Genetic and environmental influences on the size of specific brain regions in midlife the VETSA MRI study J . Neuroimage, 2010, 49 2 1213-1223.
- 141 Rimol L M, Panizzon M S, Fennema-Notestine C, et al. Cortical thickness is influenced by regionally specific genetic factors J . Biol Psychiatry, 2010, 67 5 493-499.
- 142 Stein J L, Medland S E, Vasquez A A, et al. Identification of common variants associated with human hippocampal and intracranial volumes J . Nat Genet, 2012, 44 5 552-561.
- 143 Chaste P, Klei L, Sanders S J, et al. A Genome-wide Association Study of Autism Using the Simons Simplex Collection Does Reducing Phenotypic Heterogeneity in Autism Increase Genetic Homogeneity? J . Biol Psychiatry, 2014,
- 144 Hibar D P, Stein J L, Kohannim O, et al. Voxelwise gene-wide association study vGeneWAS multivariate gene-based association testing in 731 elderly subjects J . Neuroimage, 2011, 56 4 1875-1891.
- 145 Yang J, Benyamin B, McEvoy B P, et al. Common SNPs explain a large proportion of the heritability for human height J . Nat Genet, 2010, 42 7 565-569.
- 146 Medland S E, Jahanshad N, Neale B M, et al. Whole-genome analyses of whole-brain data working within an expanded search space J . Nat Neurosci, 2014, 17 6 791-800.
- 147 International Schizophrenia C, Purcell S M, Wray N R, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder J . Nature, 2009, 460 7256 748-752.