NeuroImage 135 (2016) 186-196



Contents lists available at ScienceDirect

NeuroImage

journal homepage: www.elsevier.com/locate/ynimg

# Serotonin transporter polymorphism alters citalopram effects on human pain responses to physical pain



NeuroImage



Yina Ma <sup>a,e,f,\*</sup>, Chenbo Wang <sup>a,b</sup>, Siyang Luo <sup>a,b</sup>, Bingfeng Li <sup>b,d</sup>, Tor D. Wager <sup>g</sup>, Wenxia Zhang <sup>b,d</sup>, Yi Rao <sup>b,d</sup>, Shihui Han <sup>a,b,c,\*\*</sup>

<sup>a</sup> Department of Psychology, Peking University, Beijing, China

<sup>b</sup> PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

<sup>c</sup> Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China

<sup>d</sup> Peking-Tsinghua Center for Life Sciences at School of Life Sciences, Peking University, Beijing, China

<sup>e</sup> State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing 100875, China

<sup>f</sup> IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing 100875, China

<sup>g</sup> Psychology Department, University of Colorado Boulder, Boulder, CO, USA

#### ARTICLE INFO

Article history: Received 11 November 2015 Revised 23 April 2016 Accepted 26 April 2016 Available online 27 April 2016

Keywords: SSRI 5-HTTLPR fMRI Pain perception Personalized medicine

## ABSTRACT

Humans exhibit substantial inter-individual differences in pain perception, which contributes to variability in analgesic efficacy. Individual differences in pain sensitivity have been linked with variation in the serotonin transporter gene (5-HTTLPR), and selective serotonin reuptake inhibitors (SSRIs) such as citalopram have been increasingly used as treatments for multiple pain conditions. We combined genotyping, pharmacological challenge, and neuroimaging during painful electrical stimulation to reveal how serotonin genetics and pharmacology interact to influence pain perception and its underlying neurobiological mechanisms. In a double-blind, placebo-controlled procedure, we acutely administrated citalopram (30 mg po) to short/short (s/s) and long/ long (1/1) healthy male 5-HTTLPR homozygotes during functional MRI with painful and non-painful electrical stimulation. 5-HTTLPR genotype modulated citalopram effects on pain-related brain responses in the thalamus, cerebellum, anterior insula, midcingulate cortex and inferior frontal cortex. Specifically, citalopram significantly reduced pain-related brain responses in l/l but not in s/s homozygotes. Moreover, the interaction between 5-HTTLPR genotype and pain-related brain activity was a good predictor of the citalopram-induced reductions in pain reports. The genetic modulations of citalopram effects on brain-wide pain processing were paralleled by significant effects on the Neurological Pain Signature, a multivariate brain pattern validated to be sensitive and specific to physical pain. This work provides neurobiological mechanism by which genetic variation shapes brain responses to pain perception and treatment efficacy. These findings have important implications for the types of individuals for whom serotonergic treatments provide effective pain relief, which is critical for advancing personalized pain treatment.

© 2016 Elsevier Inc. All rights reserved.

## 1. Introduction

Pain is an unpleasant sensory and emotional experience and the most common reason people seek medical attention (Melnikova, 2010). However, pain treatment does not achieve analgesia for all individuals: one out of every 1.5 to 8.3 patients achieves effective pain relief (The 2007 Oxford league table of analgesic efficacy), implying potential individual differences in underlying mechanisms and treatment responses. Substantial inter-individual differences pervade all aspects of

http://dx.doi.org/10.1016/j.neuroimage.2016.04.064 1053-8119/© 2016 Elsevier Inc. All rights reserved.

pain responses, including subjective pain experience (Lanier, 1943; Kim et al., 2004), neural responses to painful stimulation (Coghill et al., 1999; Wager et al., 2013), and responses to pain treatment (Levine et al., 1981; Bruehl et al., 2013). Human studies implicate a significant genetic contribution to inter-individual differences in pain sensitivity (Norbury et al., 2007), chronic pain (Zondervan et al., 2005), and analgesic sensitivity (Mogil et al., 2003; Lötsch et al., 2009), which is paralleled in animal models (Mogil, 1999, 2009; Lötsch et al., 2009). These findings suggest that treatment efficacy depends in part on one's genetic makeup, providing both a challenge and an opportunity for personalized medicine. The high variability in drug efficacy across individuals signals a great need to stratify patients into groups, based on genetic and neurophysiological characteristics, that can help to determine which patient should get which treatment (Dib-Hajj and Waxman, 2014).

<sup>\*</sup> Correspondence to: Y. Ma, State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing 100875, China.

<sup>\*\*</sup> Correspondence to: S. Han, Department of Psychology, Peking University, Beijing 100080, China.

E-mail addresses: yma@bnu.edu.cn (Y. Ma), shan@pku.edu.cn (S. Han).

One key determinant of individual differences may lie in genetic variation of the serotonin transporter (5-HTT), a monoamine transporter protein that returns serotonin (5-HT) from the synaptic cleft to the presynaptic neuron. 5-HTT is thought to play a key role in nociceptive processing, as evidenced by 5-HTT knockout rodent studies and human studies (Vogel et al., 2003; Palm et al., 2008; Kupers et al., 2009, 2011; Lunn et al., 2015). For example, 5-HTT knockout mice, which are considered to be a model of lifelong selective serotonin reuptake inhibitor (SSRI) treatment (Bengel et al., 1998; Lesch and Heils, 2000), show reduced sensitivity to thermal pain (Vogel et al., 2003; Palm et al., 2008). A genetic polymorphism in the upstream promoter region of 5-HTT (5-HTTLPR), which has a short (s) and a long (l) variant, has been associated with variation in both clinical pain disorders and experimental pain (Cohen et al., 2002; Marziniak et al., 2005). The variants of the 5-HTTLPR affect the expression, transcriptional activity and function of 5-HTT, with l/l (compared to s/s) homozygotes exhibiting increased 5-HTT expression (Lesch et al., 1996). I/I homozygotes are more sensitive to experimental pain relative to s-allele carriers (Palit et al., 2011; Lindstedt et al., 2011). Moreover, serotonergic drugs, such as SSRIs, have been used for multiple pain conditions (Sindrup et al., 1992; Otto et al., 2008; Lee and Chen, 2010; Lunn et al., 2015), including poststroke pain, fibromyalgia, and neuropathic pain, though the clinical outcomes of SSRI treatment of neuropathic pain were generally modest (Finnerup et al., 2015).

These findings suggest an important relationship between 5-HTTLPR and pain, but the implications for pain treatment and underlying brain mechanisms remain unclear. Do people of different 5-HTTLPR genotypes respond differently to serotonergic treatments for pain? And, if so, what neural mechanisms underlie this interaction between genotype and treatment? To solve these issues is critical for understanding how individual differences in serotonin genetics modulate the efficacy of SSRI drug, which has important implications for the personalization of pain treatment.

The current work combined genetics, pharmacology, and neuroimaging during painful electrical stimulation to elucidate the neurobiological mechanisms through which 5-HTTLPR affects pain perception and pain treatment. Citalopram is a highly selective SSRI that selectively blocks 5-HTT activity and is associated with antinociceptive effects (Gatch et al., 1998). In a double-blind, placebo-controlled withinsubjects crossover design, we acutely administrated 30 mg citalopram, a typical dose used in previous studies (Nandam et al., 2011; Mandrioli et al., 2012; Ma et al., 2015), or placebo to s/s and l/l 5-HTTLPR homozygotes in separate sessions during functional MRI when participants anticipated and received painful and non-painful electric stimulations (Fig. 1). This pharmacogenetic neuroimaging approach allowed us to examine whether and how one's genetic makeup influenced the citalopram efficacy. Numerous neuroimaging studies have consistently shown that brain regions such as the thalamus, insula, midcingulate cortex (MCC), supplemental motor area (SMA) and primary somatosensory cortex encode the intensity of nociceptive stimuli and mediate pain sensation (Peyron et al., 2000; Apkarian et al., 2005; Tracey and Mantyh, 2007; Atlas et al., 2014). In addition, researchers have identified a specific fMRI-based multivariate pattern within and across those regions that discriminates physical pain from social pain, pain anticipation and pain recall with high sensitivity and specificity (Wager et al., 2013, >

et al., 2015), citalopram administration should produce stronger reductions in pain-related brain activity and NPS responses in l/l homozygotes. Finally, stronger brain responses to painful shock—an index of individual differences in hypersensitivity—should predict the magnitude of beneficial citalopram effects on pain. These last two points constitute two effects important for personalized medicine: (a) an interaction between genotype and treatment on pain-related brain responses, and (b) prediction from pain-related brain responses to individual differences in citalopram effects on pain within and across genetic groups, respectively.

## 2. Materials and methods

#### 2.1. Participants

Fifty-six healthy males, recruited from a pool of 901 university students genotyped for 5-HTTLPR (see below), participated in this study as paid volunteers. Two participants finished the first scanning session, but skipped the second session. Four participants were excluded due to excessive head movement. Thus the final data analyses were performed on 50 participants balanced for 5-HTTLPR genotype: 25 male s/s homozygotes (18–23 years, 19.5  $\pm$  1.7 years) and 25 male l/l homozygotes (18–23 years, 19.1  $\pm$  1.3 years). Age, education, self-esteem and anxiety trait did not differ between s/s and l/l groups (Table S1). All participants were right-handed, and had normal or corrected-to-normal vision. Exclusion criteria included any history of cardiac, hepatic, renal, pulmonary, neurological, psychiatric or gastrointestinal disorders, medication/drug use, and personal or family history of major depression or bipolar affective disorder. We tested only males in this study because we aimed to provide the first test of a complex phenomenon-a Genotype × Treatment interaction—and we wanted to avoid as many potentially confounding variables as possible in this initial test. There are documented sex differences in pain threshold (Chesterton et al., 2003; Kim et al., 2004; Fillingim et al., 2009) and pain-related brain activity (Fillingim et al., 2009; Kano et al., 2013) in the literature. Moreover, pain thresholds vary across the menstrual cycle (Riley et al., 1999; Stening et al., 2007; Fillingim et al., 2009), we aimed to test the interaction (Genotype × Treatment) of primary interest in the study without additional individual variability related to sex and menstrual phase in the initial study. This means that future studies will be required to assess the generalization of these effects to female participants.

The experimental procedures were in line with the standards set by the Declaration of Helsinki and were approved by the Research Ethics Committee of Department of Psychology, Peking University, China. Participants provided their written informed consent after all the experimental procedures had been fully explained, and were acknowledged their right to withdraw at any time during the study. Participants were compensated for their time participated in the study.

## 2.2. DNA isolation and analysis

We used established polymerase chain reaction (PCR)-based method (Ota et al., 2007) to determine the genotypes of 5-HTTLPR. In a total volume of 50  $\mu$ l, about 25 ng of genomic DNA was amplified in the presence of 1 × TransStart FastPfu DNA Polymerase (TransGen Biotech) reaction system and oligonucleotide primers (forward 5'-GCATCC CCCATTATCCCCCCT-3' and reverse 5'-AGGCTTGGAGGCCGGGATGC-3') at final concentration of 200 nM. Thermal cycling consisted of 15 min of initial denaturation at 95 °C followed by 35 cycles of 95 °C (20 s), 69 °C (20 s) and 72 °C (15 s) each with a final extension step of 10 min at 72 °C. Subsequently, PCR product was loaded onto a 3% agarose gel (BioWest G-10), to perform electrophoresis to distinguish genotypes of s/s, s/l and l/l. All genotyping was performed in duplicate.

Blood samples of 901 university students (490 males and 411 females, 18–33 years, mean age  $\pm$  SD = 19.99  $\pm$  2.76 years) were collected for 5-HTTLPR genotyping. Among 901 participants, there

were 88 long allele homozygotes (l/l), 194 heterozygotes (l/s), and 619 short allele homozygotes (s/s). The allele frequency is 79% for s-allele and 21% for l-allele, which is similar to those reported in previous studies of Asian populations (Kim et al., 2000; Tsai et al., 2002; Zhang et al., 2009; Zhang et al., 2015).

## 2.3. Stimulus and procedure

Each participant attended two sessions (at least 7 days apart, ranging 7–29 days, 14.3  $\pm$  6.7 days) and received single doses of citalopram or placebo in each session in a double-blind fully counterbalanced design (Fig. 1). To minimize potential effects of treatment order, we counterbalanced the order of citalopram/placebo treatments within each genotype group and across two groups. Moreover, we estimated the effect of treatment order by running repeated measure analyses of variance (ANOVAs) with treatment order as a between-subject variable and no significant treatment order effect was revealed.

We used a single administration of 30 mg citalopram, a typical dose (20–60 mg, Mandrioli et al., 2012; Ma, 2015) used in previous studies (Nandam et al., 2011; Ma et al., 2015). Before each fMRI session participants completed the Positive and Negative Affect Scale (PANAS, Watson et al., 1988), a 20-item self-report measure of current positive and negative affective states, and then took citalopram or placebo orally. Since pharmacokinetic studies show that citalopram is rapidly absorbed after oral administration, with plasma concentration reaching peak

100-Hz train of 0.5 ms electrical pulses with a duration of 3 s. The current intensity for 'non-painful' and 'painful' shocks was determined on an individual basis. Shocks, starting from 0.2 mA, were applied to participants and were repeated, raising 0.2 mA each time. The current intensity of the shock, to which participants answered "yes" to the question "can you feel this shock?", defined the sensory threshold. Experimenter raised the intensity and asked "is this shock painful?" Pain threshold was defined as the intensity of the shock to which participants acknowledged pain. Pain tolerance threshold was set at the maximum level of current intensity that participants could tolerate by answering "no" to the question "can you tolerate a stronger shock?" (see Table S2 for the stimuli intensity for the sensory and pain tolerance thresholds). The current intensities of sensory threshold and pain tolerance threshold were used as 'non-painful' and 'painful' stimulation during scanning, respectively (see Supplementary Methods for details about adjusted sensory and pain tolerance thresholds).

## 2.5. Imaging parameters

Functional images were acquired using 3.0-Tesla Siemens-Trio at the Beijing MRI Center for Brain Research. Blood oxygen level dependent (BOLD) gradient echo planar images were obtained using a 12-channel head coil ( $64 \times 64 \times 32$  matrix with  $3.75 \times 3.75 \times 5.0$  mm spatial resolution, repetition time (TR) = 2000 ms, echo time (TE) = 30 ms, flip angle (FA) = 90°, field of view =  $24 \times 24$  cm) during the pain task. A high-resolution T1-weighted structural image ( $256 \times 256 \times 144$  matrix with a spatial resolution of  $1 \times 1 \times 1.33$  mm,

7°) was subsequently acquired.

2.6. Imaging analysis

(GLM) for event-related designs in SPM8. The functional images were corrected for differences in acquisition time between slices for each whole-brain volume and realigned within and across runs to correct for head movement. The anatomical image was coregistered with the

were applied to functional images, which were resampled to an isotropic voxel size of  $2\times 2\times 2$  mm

'painful non-painful

The contrasts of 'painful

pain experience, respectively. In two analyses, we subjected contrast

scaling of contrast weights applied to the first-level activation parameter estimates (i.e., beta images). For a more complete discussion of variables relevant for absolute scaling of the activation parameter estimates and our estimated scaling factor, see the Supplementary Materials. Here, we reported NPS responses in units of rescaled contrast estimates, which were  $36.5 \times$  lower than the raw contrast estimates from the first-level model, though we note that the statistical comparisons we report were identical whether the rescaling is applied or not.

## 2.8. Regression analyses

We performed moderated regression analyses to examine whether 5-HTTLPR genotype moderated the relationship between the magnitude of brain responses to painful stimulation and treatment efficacy. In the moderated regression model, the independent variable (IV) was brain sensitivity to pain, defined as brain responses to painful events under placebo in an individual pain-related brain region-or, in other analyses, the NPS pattern. The dependent variable (DV) was treatment efficacy, defined here as the citalopram effect on pain reports. This effect was calculated as the differential subjective pain reports under placebo minus those under citalopram sessions. Positive values indicated that citalopram decreased pain reports, whereas negative values indicated that citalopram increased pain. The moderator was 5-HTTLPR genotype, coded as a dichotomous dummy variable in which 0 represented s/s homozygotes and 1 represented l/l homozygotes. The interactions between brain sensitivity to pain and genotype were calculated by multiplying the normalized variables together (Aiken and West, 1991). Normalized genotype, IV, and genotype  $\times$  IV interactions (moderation effects) were sequentially entered into the model. Post-hoc regression analyses were then conducted for each genotype group. This analysis identified whether brain sensitivity to pain predicted treatment efficacy, and whether its effects were moderated by genotype. If so, these variables could be used to predict who will respond to citalopram treatment, and thus personalize treatment by prospectively selecting individuals for citalopram treatment who will respond.

## 3. Results

## 3.1. Subjective pain reports and mood ratings

The pain reports were defined as the mean trial-by-trial painfulness rating scores of 'painful' stimulation and were then subjected to Treatment (citalopram vs. placebo) × Genotype (s/s vs. l/l) ANOVA. There was no significant Treatment × Genotype interaction (F(1,48) = 0.21, p = 0.65, Table 1). The mean pain reports tended to be smaller in the citalopram session than in the placebo session, similar to previous report (Gatch et al., 1998), but this effect did not reach significance (F(1,48) = 0.73, p = 0.40). Citalopram did not significantly affect pain reports in either s/s (F(1,24) = 0.10, p = 0.75) or l/l (F(1,24) = 0.69, p = 0.42) groups, motivating additional analyses of individual differences.

The post-scan rating scores of the fearful, anxious and uncomfortable feelings of electric shocks during scanning were subjected to 2 (Pain: 'painful' vs. 'non-painful')  $\times$  2 (Treatment: citalopram vs. placebo)  $\times$  2 (Genotype: s/s vs. l/l) ANOVAs. There were only significant main effects of Pain on these ratings (Fearful: F(1,48) = 407.11, p < 0.001; Anxious: F(1,48) = 378, p < 0.001; Uncomfortable: F(1,48) = 723.85, p < 0.001, Table S3). No other significant effect was observed (all p > 0.05). Self-reported mood changes from pre- to post-scan were subjected to Treatment × Genotype ANOVAs. These analyses did not show any significant effect on either positive (all p > 0.2) or negative mood (all p > 0.3, Table S4), suggesting a null effect of citalopram treatment on subjects' general mood. Therefore, the observed citalopram effects on pain processing cannot be attributed to citalopram influences on general affective states.

## 3.2. Genetic × treatment effects on neural activity during physical pain

To validate the manipulation of electric shocks at a neural level, we first calculated the contrasts of 'painful' vs. 'non-painful' shocks collapsing across s/s and l/l genotypes under placebo. Significant activations were identified using a threshold of p < 0.05 (cluster-level FDR corrected). Similar to previous findings (Peyron et al., 2000; Wager et al., 2013), under placebo, painful compared with non-painful shocks increased activity in the typical pain-related circuit, including the bilateral anterior insula (AI), posterior insula (PI), cerebellum, secondary somatosensory area (SII), thalamus (extending to basal ganglia), midbrain, MCC, SMA, and superior parietal cortex (SPC) (Fig. S1B, Fig. S2).

perception, we identified a significant Genotype × Treatment interaction on NPS responses (F(1,47) = 6.92 p = 0.012, Fig. 3A). Post hoc analyses confirmed that citalopram significantly decreased NPS responses in l/l (F(1,23) = 6.57, p = 0.018, Fig. 3) but not in s/s homozygotes (F(1,23) = 1.95, p = 0.18). These results suggested that the Genotype × Treatment interaction was also manifested at the level of a pattern of fMRI activity across multiple pain-related brain regions, consistent with the effects observed within each pain-related brain region.

3.4. Citalopram effect on pain reports: prediction from brain responses to pain reports

Our findings provided evidence for 5-HTTLPR genotype differences in the citalopram effect on pain-related activity. Given that citalopram has been increasingly used in pain treatment, it is important and of clinical interest to evaluate whether brain sensitivity to painful stimulation (measured by the magnitude of activity increases) can predict treatment efficacy, defined here as the citalopram effect on subjective pain



Fig. 2. Genotype × Treatment interaction on physical pain. Significant Genotype × Treatment interaction was observed in the bilateral thalamus, cerebellum, right Al, MCC, right inferior frontal and right lateral middle frontal (at a threshold of p < 0.05, cluster-level FDR corrected). The parameter estimates of signal intensity to 'painful' and 'non-painful' shocks were extracted from spheres with 5-mm radii centered at the peak voxel of the brain regions that was revealed in the whole-brain Genotype × Treatment interactions. Resulting means and standard errors of the mean are plotted to illustrate the contribution of each condition to the Genotype × Treatment interaction.

reports. If so, the measurement of baseline pain-related brain responses can be useful in predicting treatment efficacy and guiding treatment decisions. More importantly, we were interested in whether such prediction was moderated by the 5-HTTLPR genotype, which may also be a critical factor. Brain sensitivity to painful stimulation, genotype, and their interaction, were entered as regressors for the regression analyses of the treatment efficacy.

These analyses showed that the relationship between cerebellum/AI sensitivity to painful shocks under placebo and the treatment efficacy was significantly moderated by 5-HTTLPR genotype (right AI:

citalopram effects on pain reports. Stronger cerebellum/AI activity to painful shocks (without treatment) predicted greater citalopraminduced pain-report reduction in l/l homozygotes. However, for s/s homozygotes, citalopram treatment decreased pain reports to a greater degree in those who showed weaker cerebellum/AI activity to painful shocks. These genetic modulation effects were paralleled with significant Genotype  $\times$  Treatment interactions on the NPS — a pattern of activity across multiple brain regions associated with physical pain perception. These results indicate that one's genetic makeup interacts with baseline neural responses to pain to influence the effect of citalopram on pain perception. The current finding has important implications for patient stratification and increasing efficacy of pain treatment.

The current finding may be related to reduced descending inhibition of nociceptive signaling, which is mediated by serotonergic function in a number of animal models of descending pain control (Millan, 2002 for a systematic review). Previous studies of human participants have shown evidence for the involvement of opioidergic and dopaminergic systems in modulations of pain (Bushnell et al., 2013). However, the role for central serotonergic systems has not been well established in spite of the use of SSRIs in treating several forms of pain, such as central poststroke pain and neuropathic pain (Sindrup et al., 1992; Otto et al., 2008; Lee and Chen, 2010; Lunn et al., 2015). Our fMRI results provide the first pharmacogenetic neuroimaging evidence for the effects of both 5-HTT genotype and SSRIs as well as their interaction on the neural mediators of acute pain, demonstrating a key role of the serotonergic system in acute pain processing and providing a neural basis for further understanding of individual differences in serotonergic treatment efficacy.

The type of interactions observed here - involving a genetic background variable that predicts who will respond to drug treatment and who will not - has significance for advancing personalized medicine. There is a critical need for developing tests that can identify classes of individuals who will and will not respond to treatment, an approach referred as 'stratified medicine' (Kapur et al., 2012), which aims to increase drug efficacy. However, there have been few empirical demonstrations that individual differences in drug efficacy can be predicted (Ma et al., 2015). Regarding clinical treatment of pain, although SSRIs have been used for multiple pain conditions (Sindrup et al., 1992; Otto et al., 2008; Lee and Chen, 2010; Lunn et al., 2015), the efficacy of SSRIs was modest and treatment effects on clinical pain have been overstated in published studies (Finnerup et al., 2015). Our findings identified two strata of individuals (1/1 and s/s genotype of 5-HTTLPR) who showed differences in the efficacy of acute SSRI treatment and provide a potential account for the modest SSRI efficacy of clinical pain treatment. Furthermore, our findings indicate that not only one's genetic makeup, but also one's baseline pain responsiveness and their interactions, altered the citalopram effect on pain. We found that the pain-related brain activity predicted individual differences in treatment efficacy within and across 5-HTTLPR genotype groups, respectively. I/I homozygotes responding stronger to painful stimulation and s/s homozygotes responding weaker to painful stimulation are more likely to benefit from citalopram. It has been suggested that variations in pain characteristics between patients influence efficacy of pain treatment and patient stratification on the basis of symptoms would be beneficial on pain treatment (Dib-Hajj and Waxman, 2014). Our results suggest that genetic makeup should be taken into consideration when exploring the relationship between individuals' pain characteristic and treatment effect. Moreover, a simple genetic screen and baseline pain response measure can provide a good estimation of potential citalopram effects on pain reports. Although clinical MRI scans are not yet routinely available for this purpose, findings like these are needed to motivate their use in clinical applications.

It should be noted that the current study did not find significant 5-HTTLPR or citalopram effects on subjective pain reports. This limits the implication of the current findings in clinical populations, but does not



Fig. 4. Relationship between pain-related activity and citalopram effect on subjective painfulness ratings of painful shocks. A/B) Stronger rAI/cerebellum activity to painful shocks predicted greater decreases (for I/l homozygotes) or increases (for s/s homozygotes) in pain reports by citalopram. C) Stronger NPS responses to painful shocks under placebo predicted greater decreases in pain reports by citalopram in I/l but not in s/s homozygotes.

indicate a lack of 5-HTTLPR and citalopram influences on affective states or pain reports in general. It is possible that the effect of single doses of citalopram on subjective feeling was subtle and unable to be picked up with the current sample size. The strong pain stimulation may also lead to limited variability in pain-reports between placebo and citalopram sessions. Although there is no overall genotype/treatment effect on subjective pain reports, AI/cerebellum activity and NPS responses to pain predicted individual differences in the citalopram effect on pain reports. In addition, the pattern of neural responses and subjective reports during pain showed the same direction (i.e., I/I, relative to s/s, homozygotes showed stronger citalopram effect on neural responses and subjective reports during pain).

Pain is a multidimensional experience, involving sensorydiscriminative, affective-motivational and cognitive components (Peyron et al., 2000; Bushnell et al., 2013). The contralateral thalamus, SI/SII, and posterior insula underlie pain sensation and encode intensity of pain stimuli (Peyron et al., 2000; Bushnell et al., 1999; Ostrowsky et al., 2002; Apkarian et al., 2005). The medial thalamus, MCC, AI and cerebellum mediate pain-related affective processes of pain, for example, pain-related unpleasantness is associated with activity in the MCC, AI and cerebellum (Rainville et al., 1997; Singer et al., 2004; Craig, 2011; Bushnell et al., 2013) whereas activity increases in the thalamus reflect a general arousal reaction to pain (Peyron et al., 2000). The posterior parietal and prefrontal cortices play key roles in cognitiveevaluative aspects of pain experience (Peyron et al., 2000; Bushnell et al., 1999). To date, little is known about whether the activities in different regions of the pain-related network are similarly modulated by 5-HTT. Our findings revealed that the interaction between serotonergic genotype and drug influenced the thalamus, AI, MCC, cerebellum and SMA activity but did not produce significant effects on the SI, SII, PI and parietal activity to pain, suggesting that 5-HTT plays a more important role in enhancing motor-related and affective (relative to sensory or cognitive) processes during pain experience.

In line with this result, we found that the interaction between 5-HTTLPR genotype and individual's baseline neural response to pain was a good predictor of the potential analgesic effect of citalopram. Stronger baseline cerebellum/AI activity to painful shocks (without treatment) predicted greater citalopram-induced subjective painreport reduction in l/l homozygotes. This effect was mainly observed in the AI, cerebellum, and MCC (to a less extent, as the interaction between 5-HTTLPR genotype and baseline MCC activity only marginally predicted citalopram effect on pain reports,  $\beta = 0.29$ , p = 0.072). These regions are implicated in the affective and motor-related component of pain (Peyron et al., 2000; Moulton et al., 2010; Bushnell et al., 2013). Expectation of pain activated the medial frontal, AI, MCC and cerebellum (Ploghaus et al., 1999; Chua et al., 1999), and observing another individual in pain activated AI, MCC, cerebellum and brainstem (Singer et al., 2004: Jackson et al., 2005: Xu et al., 2009). Moreover, these regions are particularly important for the subjective experience of pain, encoding subjectively perceived pain intensity (Peyron et al., 2000; Koyama et al., 2005; Baliki et al., 2009) and are sensitive to interindividual differences (Coghill et al., 2003; Coghill, 2010). Consistently, we showed that interaction between 5-HTTLPR and baseline NPS values predicted citalopram effect on pain report. Given that the NPS pattern mainly reflects sensory and affective aspects of pain (Wager et al., 2013; Woo et al., 2015), these findings together suggest that neural activity in the affective node of the pain network provides a good prediction of individual differences in citalopram effect on subjective pain reports, with the prediction directions depending on the 5-HTTLPR genotype.

The brain activity pattern observed in our work during pain anticipation was different from the previous report of greater neural activity in response to acute stress induced by pain anticipation in s/s than l allele carriers of 5-HTTLPR (Drabant et al., 2012), which was apparently different from our results of absence of genotype differences in brain activity during pain anticipation and the stronger neural activity during experiencing pain in 1/l compared to s/s homozygotes. There were several factors that might induce the difference in fMRI results across the studies. First, it has been recognized that, although a similar neural network is shared by anticipation of pain and experience of pain, the neural activity activated during experiencing pain and anticipation of pain can show distinct patterns in the same brain region (e.g., insular cortex and cerebellum, Ploghaus et al., 1999). Similar to Drabant et al. (2012), our previous studies found that s/s compared to l/l homozygotes exhibit greater neural responses to threatening signals such as fearful faces (Ma et al., 2015) and reflection on one's own undesirable personality traits (Ma et al., 2014a). In contrast, l/l homozygotes compared to sallele carriers are more sensitive to pain stimulation (Palit et al., 2011; Lindstedt et al., 2011) and showed hyperactivity within the painrelated network when experiencing physical pain (current work). Therefore, it is likely that 5-HTTLPR may produce distinct effects on neural activity in response to acute stress induced by anticipation and experience of physical pain. Second, there was a key difference in the design of anticipation of pain between Drabant et al. (2012) and the current work. In our study, an electrical shock followed each cue. This design did not engage uncertainty of pain stimulations and thus might reduce the fear and stress related to pain stimulation. In order to maximally inlong been recognized (Moulton et al., 2010), evidenced by the cooccurrence of pain condition and depression (Katona et al., 2005) and the common brain regions modulated by pain and depression (Chopra and Arora, 2014). However, the experience of pain is different from the fear and anxiety caused by threats of pain (Ploghaus et al., 1999). Future research should further clarify distinct SSRI effects on pain conditions and negative emotion in order to predict SSRI effects in clinical treatment.

In conclusion, this study elucidates the neurobiological mechanisms underlying the serotonergic genetics modulation of SSRI-induced brain changes during physical pain perception. As SSRIs have been increasingly used for multiple pain conditions (Sindrup et al., 1992; Otto et al., 2008; Lee and Chen, 2010; Lunn et al., 2015), the current finding of serotonin genetics and pharmacology interaction has implications for the types of individuals for whom serotonergic treatments may provide effective pain relief. Although our experimental data from healthy volunteers have implications for clinical practice, the generalizability of our findings must be assessed on chronic pain conditions and patient population. Future research should examine the role of 5-HTT and SSRI treatment in patients with various chronic pain conditions, thus to promote personalized pain treatment. Finally, the current finding was observed from a sample of only males, future research should also examine whether the current findings can be generalized to females.

## Conflict of interest

The author declares no conflict of interest.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (31421003, 31470986, 91332125, 81161120539), the National Basic Research Program of China (973 Program 2010CB833901, 2010CB833903), Beijing Municipal Natural Science Foundation (Z111107067311058), and the Ministry of Education of China (20130001110049). We thank Y Liu, Y Zhang, Z Shi, X Zuo, D Yu, R Sun, X Li for their help with participant recruitment.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at http://dx. doi.org/10.1016/j.neuroimage.2016.04.064.

#### Reference

- Aiken, L.S., West, S.G., 1991. Multiple Regression: Testing and Interpreting Interactions. Sage, Newbury Park, CA.
- Apkarian, A.V., Bushnell, M.C., Treede, R.D., Zubieta, J.K., 2005. Human brain mechanisms of pain perception and regulation in health and disease. Eur. J. Pain 9, 463–484.
- Atlas, LY., Lindquist, M.A., Bolger, N., Wager, T.D., 2014. Brain mediators of the effects of noxious heat on pain. Pain 155, 1632–1648.
- Baliki, M.N., Geha, P.Y., Apkarian, A.V., 2009. Parsing pain perception between nociceptive representation and magnitude estimation. J. Neurophysiol. 101, 875–887.
- Bandolier Extra (Evidence-based health care). Acute pain. The 2007 Oxford league table of analgesic efficacy. Bandolier. Available from: http://www.medicine.ox.ac.uk/ bandolier/booth/painpag/acutrev/analgesics/leagtab.html.
- Bengel, D., Murphy, D.L., Andrews, A.M., Wichems, C.H., Feltner, D., Heils, A., Mössner, R., Westphal, H., Lesch, K.P., 1998. Altered brain serotonin homeostasis and locomotor insensitivity to 3, 4-methylenedioxymethamphetamine ("ecstasy") in serotonin transporter-deficient mice. Mol. Pharmacol. 53, 649–655.
- Bruehl, S., Apkarian, A.V., Ballantyne, J.C., Berger, A., Borsook, D., Chen, W.G., Farrar, J.T., Haythornthwaite, J.A., Horn, S.D., Iadarola, M.J., Inturrisi, C.E., 2013. Personalized medicine and opioid analgesic prescribing for chronic pain: opportunities and challenges. J. Pain 14, 103–113.
- Bushnell, M.C., Duncan, G.H., Hofbauer, R.K., Ha, B., Chen, J.I., Carrier, B., 1999. Pain perception: is there a role for primary somatosensory cortex? Proc. Natl. Acad. Sci. U. S. A. 96, 7705–7709.
- Bushnell, M.C., Ceko, M., Low, L.A., 2013. Cognitive and emotional control of pain and its disruption in chronic pain. Nat. Rev. Neurosci. 14, 502–511.
- Carlsson, K., Andersson, J., Petrovic, P., Petersson, K.M., Ohman, A., Ingvar, M., 2006. Predictability modulates the affective and sensory-discriminative neural processing of pain. Neuroimage 32, 1804–1814.

- Chesterton, L.S., Barlas, P., Foster, N.E., Baxter, G.D., Wright, C.C., 2003. Gender differences in pressure pain threshold in healthy humans. Pain 101, 259–266.
- Chopra, K., Arora, V., 2014. An intricate relationship between pain and depression: clinical correlates, coactivation factors and therapeutic targets. Expert Opin. Ther. Targets 18, 159–176.
- Chua, P., Krams, M., Toni, I., Passingham, R., Dolan, R., 1999. A functional anatomy of anticipatory anxiety. NeuroImage 9, 563–571.
   Coghill, R.C., 2010. Individual differences in the subjective experience of pain: new in-
- Coghill, R.C., 2010. Individual differences in the subjective experience of pain: new insights into mechanisms and models. Headache 50, 1531–1535.
- Coghill, R.C., Sang, C.N., Maisog, J.M., Iadarola, M.J., 1999. Pain intensity processing within the human brain: a bilateral, distributed mechanism. J. Neurophysiol. 82, 1934–1943.
- Coghill, R.C., McHaffie, J.G., Yen, Y.F., 2003. Neural correlates of interindividual differences in the subjective experience of pain. Proc. Natl. Acad. Sci. U. S. A. 100, 8538–8542.
- Cohen, H., Buskila, D., Neumann, L., Ebstein, R.P., 2002. Confirmation of an association between fibromyalgia and serotonin transporter promoter region (5-HTTLPR) polymorphism, and relationship to anxiety-related personality traits. Arthritis Rheum. 46, 845–847.
- Craig, A.D., 2011. Significance of the insula for the evolution of human awareness of feelings from the body. Ann. N. Y. Acad. Sci. 1225, 72–82.
- Dib-Hajj, S.D., Waxman, S.G., 2014. Translational pain research: lessons from genetics and genomics. Sci. Transl. Med. 6, 249sr4.
- Drabant, E.M., Ramel, W., Edge, M.D., Hyde, L.W., Kuo, J.R., Goldin, P.R., Hariri, A.R., Gross, J.J., 2012. Neural mechanisms underlying 5-HTTLPR-related sensitivity to acute stress. Am. J. Psychiatry 169, 397–405.
- Fillingim, R.B., King, C.D., Ribeiro-Dasilva, M.C., Rahim-Williams, B., Riley, J.L., 2009. Sex, gender, and pain: a review of recent clinical and experimental findings. J. Pain 10, 447–485.
- Finnerup, N.B., Attal, N., Haroutounian, S., McNicol, E., Baron, R., Dworkin, R.H., Gilron, I., Haanpää, M., Hansson, P., Jensen, T.S., Kamerman, P.R., 2015. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. Lancet Neurol. 14, 162–173.
- Gatch, M.B., Negus, S.S., Mello, N.K., 1998. Antinociceptive effects of monoamine reuptake inhibitors administered alone or in combination with mu opioid agonists in rhesus monkeys. Psychopharmacology 135, 99–106.
- Hariri, A.R., Mattay, V.S., Tessitore, A., Kolachana, B., Fera, F., Goldman, D., Egan, M.F., Weinberger, D.R., 2002. Serotonin transporter genetic variation and the response of the human amygdala. Science 297, 400–403.
- Hu, X.Z., Rush, A.J., Charney, D., Wilson, A.F., Sorant, A.J., Papanicolaou, G.J., Fava, M., Trivedi, M.H., Wisniewski, S.R., Laje, G., Paddock, S., 2007. Association between a functional serotonin transporter promoter polymorphism and citalopram treatment in adult outpatients with major depression. Arch. Gen. Psychiatry 64, 783–792.
- Jackson, P.L., Meltzoff, A.N., Decety, J., 2005. How do we perceive the pain of others? A window into the neural processes involved in empathy. NeuroImage 24, 771–779.
- Kano, M., Farmer, A.D., Aziz, Q., Giampietro, V.P., Brammer, M.J., Williams, S.C., Fukudo, S., Coen, S.J., 2013. Sex differences in brain response to anticipated and experienced visceral pain in healthy subjects. Am. J. Physiol. Gastrointest. Liver Physio. 304, G687–G699.
- Kapur, S., Phillips, A.G., Insel, T.R., 2012. Why has it taken so long for biological psychiatry to develop clinical tests and what to do about it? Mol. Psychiatry 17, 1174–1179.
- Katona, C., Peveler, R., Dowrick, C., Wessely, S., Feinmann, C., Gask, L., Lloyd, H., de C Williams, A.C., Wager, E., 2005. Pain symptoms in depression: definition and clinical significance. Clin. Med. 5, 390–395.
- Kim, D.K., Lim, S.W., Lee, S., Sohn, S.E., Kim, S., Hahn, C.G., Carroll, B.J., 2000. Serotonin transporter gene polymorphism and antidepressant response. Neuroreport 11, 215–219.
- Kim, H., Neubert, J.K., San Miguel, A., Xu, K., Krishnaraju, R.K., Iadarola, M.J., Goldman, D., Dionne, R.A., 2004. Genetic influence on variability in human acute experimental pain sensitivity associated with gender, ethnicity and psychological temperament. Pain 109, 488–496.
- Koyama, T., McHaffie, J.G., Laurienti, P.J., Coghill, R.C., 2005. The subjective experience of pain: where expectations become reality. Proc. Natl. Acad. Sci. U. S. A. 102, 12950–12955.
- Kupers, R., Frokjaer, V.G., Naert, A., Christensen, R., Budtz-Joergensen, E., Kehlet, H., Knudsen, G.M., 2009. A PET [18F]altanserin study of 5-HT2A receptor binding in the human brain and responses to painful heat stimulation. NeuroImage 44, 1001–1007.
- Kupers, R., Frokjaer, V.G., Erritzoe, D., Naert, A., Budtz-Joergensen, E., Nielsen, F.A., Kehlet, H., Knudsen, G.M., 2011. Serotonin transporter binding in the hypothalamus correlates negatively with tonic heat pain ratings in healthy subjects: a [11C]DASB PET study. NeuroImage 54, 1336–1343.
- Lanier, L.H., 1943. Variability in the pain threshold. Science 97, 49-50.
- Lee, Y.C., Chen, P.P., 2010. A review of SSRIs and SNRIs in neuropathic pain. Expert. Opin. Pharmacother. 11, 2813–2825.
- Lee, B.T., Ham, B.J., 2008. Serotonergic genes and amygdala activity in response to negative affective facial stimuli in Korean women. Genes Brain Behav. 7, 899–905.
  Lesch, K.P., Heils, A., 2000. Serotonergic gene transcriptional control regions: targets for
- antidepressant drug development? Int. J. Neuropsychopharmacol. 3, 67–79.
  Lesch, K.P., Bengel, D., Heils, A., Sabol, S.Z., Greenberg, B.D., Petri, S., Benjamin, J., Müller,
- Lestri, K.F., Benger, D., Hens, A., Sabol, S.Z., Greenberg, D.J., Petri, S., Benjarim, J., Muner, C.R., Hamer, D.H., Murphy, D.L., 1996. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. Science 274, 1527–1531.
- Levine, J.D., Gordon, N.C., Smith, R., Fields, H.L., 1981. Analgesic responses to morphine and placebo in individuals with postoperative pain. Pain 10, 379–389.
- Lindstedt, F., Lonsdorf, T.B., Schalling, M., Kosek, E., Ingvar, M., 2011. Perception of thermal pain and the thermal grill illusion is associated with polymorphisms in the serotonin transporter gene. PLoS One 6, e17752.

- Long, H., Liu, B., Hou, B., Wang, C., Li, J., Qin, W., Wang, D., Zhou, Y., Kendrick, K., Yu, C., Jiang, T., 2013. The long rather than the short allele of 5-HTTLPR predisposes Han Chinese to anxiety and reduced connectivity between prefrontal cortex and amygdala. Neurosci. Bull. 29, 4–15.
- Lötsch, J., Geisslinger, G., Tegeder, I., 2009. Genetic modulation of the pharmacological treatment of pain. Pharmacol. Ther. 124, 168–184.
- Lunn, T.H., Frokjaer, V.G., Hansen, T.B., Kristensen, P.W., Lind, T., Kehlet, H., 2015. Analgesic effect of perioperative escitalopram in high pain catastrophizing patients after total knee arthroplasty: a randomized, double-blind, placebo-controlled trial. Anesthesiology 122, 884–894.
- Ma, Y., 2015. Neuropsychological mechanism underlying antidepressant effect: a systematic meta-analysis. Mol. Psychiatry 20, 311–319.
- Ma, Y., Li, B., Wang, C., Shi, Z., Sun, Y., Sheng, F., Zhang, Y., Zhang, W., Rao, Y., Han, S., 2014a. 5-HTTLPR polymorphism modulates neural mechanisms of negative selfreflection. Cereb. Cortex 24, 2421–2429.
- Ma, Y., Wang, C., Li, B., Zhang, W., Rao, Y., Han, S., 2014b. Does self-construal predict activity in the social brain network? A genetic moderation effect. Soc. Cogn. Affect. Neurosci. 9, 1360–1367.
- Ma, Y., Li, B., Zhang, W., Rao, Y., Han, S., 2015. Allelic variation in 5-HTTLPR and the effects of citalopram on the emotional neural network. Br. J. Psychiatry 206, 385–392.
- Mandrioli, R., Mercolini, L., Saracino, M.A., Raggi, M.A., 2012. Selective serotonin reuptake inhibitors (SSRIs): therapeutic drug monitoring and pharmacological interactions. Curr. Med. Chem. 19, 1846–1863.
- Marziniak, M., Mössner, R., Schmitt, A., Lesch, K.P., Sommer, C., 2005. A functional serotonin transporter gene polymorphism is associated with migraine with aura. Neurology 64, 157–159.
- Melnikova, I., 2010. Pain market. Nat. Rev. Drug Discov. 9, 589–590.
- Millan, M.J., 2002. Descending control of pain. Prog. Neurobiol. 66, 355–474.
- Mogil, J.S., 1999. The genetic mediation of individual differences in sensitivity to pain and its inhibition. Proc. Natl. Acad. Sci. U. S. A. 96, 7744–7751.
- Mogil, J.S., 2009. Animal models of pain: progress and challenges. Nat. Rev. Neurosci. 10, 283–294.
- Mogil, J.S., Wilson, S.G., Chesler, E.J., Rankin, A.L., Nemmani, K.V., Lariviere, W.R., Groce, M.K., Wallace, M.R., Kaplan, L., Staud, R., Ness, T.J., 2003. The melanocortin-1 receptor gene mediates female-specific mechanisms of analgesia in mice and humans. Proc. Natl. Acad. Sci. U. S. A. 100, 4867–4872.
- Monat, A., Averill, J.R., Lazarus, R.S., 1972. Anticipatory stress and coping reactions under various conditions of uncertainty. J. Pers. Soc. Psychol. 24, 237–253.
   Moulton, E.A., Schmahmann, J.D., Becerra, L., Borsook, D., 2010. The cerebellum and pain:
- Moulton, E.A., Schmahmann, J.D., Becerra, L., Borsook, D., 2010. The cerebellum and pain: passive integrator or active participator? Brain Res. Rev. 65, 14–27.
- Nandam, L.S., Hester, R., Wagner, J., Cummins, T.D., Garner, K., Dean, A.J., Kim, B.N., Nathan, P.J., Mattingley, J.B., Bellgrove, M.A., 2011. Methylphenidate but not atomoxetine or citalopram modulates inhibitory control and response time variability. Biol. Psychiatry 69, 902–904.
- Norbury, T.A., MacGregor, A.J., Urwin, J., Spector, T.D., McMahon, S.B., 2007. Heritability of responses to painful stimuli in women: a classical twin study. Brain 130, 3041–3049.
- Ostrowsky, K., Magnin, M., Ryvlin, P., Isnard, J., Guenot, M., Mauguiere, F., 2002. Representation of pain and somatic sensation in the human insula: a study of responses to direct electrical cortical stimulation. Cereb. Cortex 12, 376–385.
- Ota, M., Fukushima, H., Kulski, J.K., Inoko, H., 2007. Single nucleotide polymorphism detection by polymerase chain reaction-restriction fragment length polymorphism. Nat. Protoc. 2, 2857–2864.
- Otto, M., Bach, F.W., Jensen, T.S., Brøsen, K., Sindrup, S.H., 2008. Escitalopram in painful polyneuropathy: a randomized, placebo-controlled, cross-over trial. Pain 139, 275–283.
- Palit, S., Sheaff, R.J., France, C.R., McGlone, S.T., Potter, W.T., Harkness, A.R., McNulty, J.L., Bartley, E.J., Hoffmann, R., Monda, J.K., Rhudy, J.L., 2011. Serotonin transporter gene (5-HTTLPR) polymorphisms are associated with emotional modulation of pain but not emotional modulation of spinal nociception. Biol. Psychol. 86, 360–369.
- Palm, F., Mössner, R., Chen, Y., He, L., Gerlach, M., Bischofs, S., Riederer, P., Lesch, K.P., Sommer, C., 2008. Reduced thermal hyperalgesia and enhanced peripheral nerve injury after hind paw inflammation in mice lacking the serotonin-transporter. Eur. J. Pain 12, 790–797.

- Peyron, R., Laurent, B., García-Larrea, L., 2000. Functional imaging of brain responses to pain. A review and meta-analysis (2000). Neurophysiol. Clin. 30, 263–288.
- Pezawas, L., Meyer-Lindenberg, A., Drabant, E.M., Verchinski, B.A., Munoz, K.E., Kolachana, B.S., Egan, M.F., Mattay, V.S., Hariri, A.R., Weinberger, D.R., 2005. 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. Nat. Neurosci. 8, 828–834.
- Ploghaus, A., Tracey, I., Gati, J.S., Clare, S., Menon, R.S., Matthews, P.M., Rawlins, J.N., 1999. Dissociating pain from its anticipation in the human brain. Science 284, 1979–1981. Rainville, P., Duncan, G.H., Price, D.D., Carrier, B., Bushnell, M.C., 1997. Pain affect encoded
- in human anterior cingulate but not somatosensory cortex. Science 277, 968–971. Riley, J.L.I., Robinson, M.E., Wise, E.A., Price, D.D., 1999. A meta-analytic review of pain
- perception across the menstrual cycle. Pain 81, 225–235.
- Rocha, A., Marques, M.P., Coelho, E.B., Lanchote, V.L., 2007. Enantioselective analysis of citalopram and demethylcitalopram in human and rat plasma by chiral LC-MS/MS: application to pharmacokinetics. Chirality 19, 793–801.
- Serretti, A., Kato, M., De Ronchi, D., Kinoshita, T., 2007. Meta-analysis of serotonin transporter gene promoter polymorphism (5-HTTLPR) association with selective serotonin reuptake inhibitor efficacy in depressed patients. Mol. Psychiatry 12, 247–257.
- Sindrup, S.H., Bjerre, U., Dejgaard, A., Brøsen, K., Aaes-Jørgensen, T., Gram, L.F., 1992. The selective serotonin reuptake inhibitor citalopram relieves the symptoms of diabetic neuropathy. Clin. Pharmacol. Ther. 52, 547–552.
- Singer, T., Seymour, B., O'Doherty, J., Kaube, H., Dolan, R.J., Frith, C.D., 2004. Empathy for pain involves the affective but not sensory components of pain. Science 303, 1157–1162.
- Stening, K., Eriksson, O., Wahren, L., Berg, G., Hammar, M., Blomqvist, A., 2007. Pain sensations to the cold pressor test in normally menstruating women: comparison with men and relation to menstrual phase and serum sex steroid levels. Am. J. Phys. Regul. Integr. Comp. Phys. 293, R1711–R1716.
- Tracey, I., Mantyh, P.W., 2007. The cerebral signature for pain perception and its modulation. Neuron 55, 377–391.
- Tsai, S., Hong, C., Cheng, C., 2002. Serotonin transporter genetic polymorphisms and harm avoidance in the Chinese. Psychiatr. Genet. 12, 165–168.
- Vogel, C., Mössner, R., Gerlach, M., Heinemann, T., Murphy, D.L., Riederer, P., Lesch, K.P., Sommer, C., 2003. Absence of thermal hyperalgesia in serotonin transporter-deficient mice. J. Neurosci. 23, 708–715.
- Wager, T.D., Rilling, J.K., Smith, E.E., Sokolik, A., Casey, K.L., Davidson, R.J., Kosslyn, S.M., Rose, R.M., Cohen, J.D., 2004. Placebo-induced changes in FMRI in the anticipation and experience of pain. Science 303, 1162–1167.
- Wager, T.D., Atlas, L.Y., Lindquist, M.A., Roy, M., Woo, C.W., Kross, E., 2013. An fMRI-based neurologic signature of physical pain. N. Engl. J. Med. 368, 1388–1397.
- Watson, D., Clark, LA., Tellegen, A., 1988. Development and validation of brief measures of positive and negative affect: the PANAS scales. J. Pers. Soc. Psychol. 54, 1063–1070.
- Whale, R., Quested, D.J., Laver, D., Harrison, P.J., Cowen, P.J., 2000. Serotonin transporter (5-HITT) promoter genotype may influence the prolactin response to clomipramine. Psychopharmacology 150, 120–122.
- Woo, C.W., Roy, M., Buhle, J.T., Wager, T.D., 2015. Distinct brain systems mediate the effects of nociceptive input and self-regulation on pain. PLoS Biol. 13, e1002036.
- Xu, X., Zuo, X., Wang, X., Han, S., 2009. Do you feel my pain? Racial group membership modulates empathic neural responses. J. Neurosci. 29, 8525–8529.
- Zhang, K., Xu, Q., Xu, Y., Yang, H., Luo, J., Sun, Y., Sun, N., Wang, S., Shen, Y., 2009. The combined effects of the 5-HTTLPR and 5-HTR1A genes modulates the relationship between negative life events and major depressive disorder in a Chinese population. J. Affect. Disord. 114, 224–231.
- Zhang, L., Liu, L., Li, X., Song, Y., Liu, J., 2015. Serotonin transporter gene polymorphism (5-HTTLPR) influences trait anxiety by modulating the functional connectivity between the amygdala and insula in Han Chinese males. Hum. Brain Mapp. http://dx.doi.org/ 10.1002/hbm.22803.
- Zondervan, K.T., Cardon, L.R., Kennedy, S.H., Martin, N.G., Treloar, S.A., 2005. Multivariate genetic analysis of chronic pelvic pain and associated phenotypes. Behav. Genet. 35, 177–188.