
Received 2012-04-30 Accepted 2012-11-23

This review was supported by the National Natural Science Foundation of China (No. 30950030), the National Basic Research Development Program (973 Program) of China (No. 2009CB320901), the Doctoral Fund of Ministry of Education of China (No. 200900011110050), and “985” grants from Peking University for Physiological Psychology.

*Corresponding author. Tel: +86-10-62756804; Fax: +86-10-62761081; E-mail: liangli@pku.edu.cn

([11–13])

[1–3] ()

() [14]

(prepulse inhibition, PPI) [4, 5] [6, 7] [15, 16] PPI ()

() [17–20]

[8] [4, 21, 22] PPI [6, 21]

PPI PPI

[3] PPI

早期社会隔离引起的认知行为异常 [3] (auditory fear conditioning) PPI

运动增多 1

(open [21] field test) [4, 9, 10] PPI [22]

[10] PPI PPI

[2, 5] Wistar Lister Hooded PPI

[1, 5] Sprague-Dawley PPI [23]

([2]) “ - ” [24] PPI

1.

Table 1. Main behavioral changes associated with social isolation of rats

Behavior	Behavioral changes associated with isolation rearing
Hyperactivity	Hyperactivity in isolation reared rats [1, 4, 5, 10–12] due to less time spent in resting while not due to velocity change [10] Handling or environment habituation could reverse this hyperactivity in isolated rats [2] Hyperactivity appears after 2 weeks of isolation [4, 11]
Startle and PPI	Baseline of startle increases [4, 21, 22] PPI decreases [6, 21] Enhancement of PPI induced by auditory fear conditioning declines [21, 22] Enhancement of PPI induced by perceived spatial separation declines [22] Possible brain areas involved are the amygdala, the posterior parietal cortex, the primary auditory cortex, and the prefrontal cortex [66]
Learning and memory	Deficit in fear conditioning [1, 21] Improvement in reward learning [29, 30, 67, 68] Deficit in reversal learning and no effect on acquisition learning [32, 33] Impairment in recognition [33]

PPI: prepulse inhibition.

[25]

[26, 27]

[27, 28]

学习和记忆功能异常

PPI

PPI

[21]

(contextual fear conditioning)

[1]

[29]

C57BL/6J

[30]

Wistar

[31]

8

[32]

Bianchi

[33]

早期

([48])
谷氨酸的异常

NMDA

[49] Melendez


mGluR1 mGluR5

[50] 26

mGluR6 AMPA3

[10]

早期社会隔离引起的



[60]

PPI

[61]

[37]

[62]

[63]

[64]

[65]

- 12 Einon DF, Morgan M. A critical period for social isolation in the rat. *Dev Psychobiol* 1977; 10(2): 123–132.
- 13 Hickey AJ, Reynolds JN, Beninger RJ. Post-weaning social isolation and subchronic NMDA glutamate receptor blockade: Effects on locomotor activity and GABA signaling in the rat suggest independent mechanisms. *Pharmacol Biochem Behav* 2012; 101(2): 231–238.
- 14 Landis C, Hunt W. *The Startle Pattern*. Oxford, England: Farrar & Rinehart, 1939.
- 15 Li L, Frost BJ. Azimuthal sensitivity of rat pinna reflex: EMG recordings from cervicoauricular muscles. *Hear Res* 1996; 100(1–2): 192–200.
- 16 Li L, Yeomans J. Summation between acoustic and trigeminal stimuli evoking startle. *Neuroscience* 1999; 90(1): 139–152.
- 17 Buckland G, Buckland J, Jamieson C, Ison J. Inhibition of startle response to acoustic stimulation produced by visual prestimulation. *J Comp Physiol Psychol* 1969; 67(4): 493–496.
- 18 Ison JR, Hammond GR. Modification of the startle reflex in the rat by changes in the auditory and visual environments. *J Comp Physiol Psychol* 1971; 75(3): 435–452.
- 19 Hoffman HS, Searle JL. Acoustic variables in the modification of startle reaction in the rat. *J Comp Physiol Psychol* 1965; 60(1): 53–58.
- 20 Pinckney LA. Inhibition of the startle reflex in the rat by prior tactile stimulation. *Learn Behav* 1976; 4(4): 467–472.
- 21 Li N, Ping J, Wu R, Wang C, Wu X, Li L. Auditory fear conditioning modulates prepulse inhibition in socially reared rats and isolation-reared rats. *Behav Neurosci* 2008; 122(1): 107–118.
- 22 Du Y, Wu X, Li L. Emotional learning enhances stimulus-specific top-down modulation of sensorimotor gating in socially reared rats but not isolation-reared rats. *Behav Brain Res* 2010; 206(2): 192–201.
- 23 Hazlett EA, Romero MJ, Haznedar MM, New AS, Goldstein KE, Newmark RE, Siever LJ, Buchsbaum MS. Deficient attentional modulation of startle eyeblink is associated with symptom severity in the schizophrenia spectrum. *Schizophr Res* 2007; 93(1–3): 288–295.
- 24 Graham F. The more or less startling effects of weak pre-stimulation. *Psychophysiology* 1975; 12: 238–248.
- 25 Li L (), Shao F. Impaired auditory sensorimotor gating: An animal model of schizophrenia. *Chin Sci Bull ()* 2003; 48(19): 2031–2037 (Chinese, English abstract).
- 26 Koch M, Schnitzler HU. The acoustic startle response in rats--circuits mediating evocation, inhibition and potentiation. *Behav Brain Res* 1997; 89(1–2): 35–49.
- 27 Braff DL, Geyer MA, Swerdlow NR. Human studies of prepulse inhibition of startle, normal subjects, patient groups, and pharmacological studies. *Psychopharmacology (Berl)* 2001; 156: 234–258.
- 28 Geyer MA, Krebs-Thomson K, Braff DL, Swerdlow NR. Pharmacological studies of prepulse inhibition models of sensorimotor gating deficits in schizophrenia: a decade in review. *Psychopharmacology (Berl)* 2001; 156(2): 117–154.
- 29 Jones G, Marsden C, Robbins TW. Increased sensitivity to amphetamine and reward-related stimuli following social isolation in rats: possible disruption of dopamine-dependent mechanisms of the nucleus accumbens. *Psychopharmacology (Berl)* 1990; 102(3): 364–372.
- 30 Advani T, Hensler JG, Koek W. Effect of early rearing conditions on alcohol drinking and 5-HT1A receptor function in C57BL/6J mice. *Int J Neuropsychopharmacol* 2007; 10(5): 595–608.
- 31 Hall F, Huang S, Fong G, Pert A, Linnoila M. Effects of isolation-rearing on voluntary consumption of ethanol, sucrose and saccharin solutions in Fawn Hooded and Wistar rats. *Psychopharmacology (Berl)* 1998; 139(3): 210–216.
- 32 Li N, Wu X, Li L. Chronic administration of clozapine alleviates reversal-learning impairment in isolation-reared rats. *Behav Pharmacol* 2007; 18(2): 135–145.
- 33 Bianchi M, Fone K, Azmi N, Heidbreder C, Hagan J, Marsden C. Isolation rearing induces recognition memory deficits accompanied by cytoskeletal alterations in rat hippocampus. *Eur J Neurosci* 2006; 24(10): 2894–2902.
- 34 Day-Wilson K, Jones D, Southam E, Cilia J, Totterdell S. Medial prefrontal cortex volume loss in rats with isolation rearing-induced deficits in prepulse inhibition of acoustic startle. *Neuroscience* 2006; 141(3): 1113–1121.
- 35 Varty GB, Marsden CA, Higgins GA. Reduced synaptophysin immunoreactivity in the dentate gyrus of prepulse inhibition-impaired isolation-reared rats. *Brain Res* 1999; 824(2): 197–203.
- 36 Silva-Gómez AB, Rojas D, Juárez I, Flores G. Decreased dendritic spine density on prefrontal cortical and hippocampal pyramidal neurons in postweaning social isolation rats. *Brain Res* 2003; 983(1–2): 128–136.
- 37 Lapid M, Fulford A, Muchimapura S, Mason R, Parker T, Marsden C. Influence of postweaning social isolation in the rat on brain development, conditioned behavior, and neurotransmission. *Neurosci Behav Physiol* 2003; 33(1): 13–29.
- 38 Djouma E, Card K, Lodge DJ, Lawrence AJ. The CRF1 receptor antagonist, antalarmin, reverses isolation-induced up-regulation of dopamine D2 receptors in the amygdala and nucleus accumbens of fawn-hooded rats. *Eur J Neurosci* 2006; 23(12): 3319–3327.
- 39 Howes OD, Kapur S. The dopamine hypothesis of schizo-

phrenia: version III—the final common pathway. Schizophr
Bull 2009;(1)Tj()Tu

- 65 Marcotte ER, Pearson DM, Srivastava LK. Animal models of schizophrenia: a critical review. *J Psychiatry Neurosci* 2001; 26(5): 395–410.
- 66 Du Y, Wu X, Li L. Differentially organized top-down modulation of prepulse inhibition of startle. *J Neurosci* 2011; 31(38): 13644–13653.
- 67 Hall F, Humby T, Wilkinson L, Robbins T. The effects of isolation-rearing on sucrose consumption in rats. *Physiol Behav* 1997; 62(2): 291–297.
- 68 Hall F, Huang S, Fong G, Pert A, Linnoila M. Effects of isolation-rearing on locomotion, anxiety and responses to ethanol in Fawn Hooded and Wistar rats. *Psychopharmacology (Berl)* 1998; 139(3): 203–209.