

Intention Modulates the Effect of Punishment Threat in Norm Enforcement via the Lateral Orbitofrontal Cortex

Yuan Zhang^{1,*}, Hongbo Yu^{1,*}, Yunlu Yin¹, and Xiaolin Zhou^{1,3,4,5}

¹Center for Brain and Cognitive Sciences and Department of Psychology, Peking University, Beijing 100871, China, ²Institute for Neurotechnology (SINAPSE), Center for Life Sciences, National University of Singapore, Singapore 117456, Singapore, ³Key Laboratory of Machine Perception (Ministry of Education), Beijing Key Laboratory of Behavior and Mental Health, and ⁴PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing 100871, China

Although economic theories suggest that punishment threat is crucial for maintaining social norms, counterexamples are noted in which punishment threat hinders norm compliance. Such discrepancy may arise from the intention behind the threat: unintentionally introduced punishment threat facilitates, whereas intentionally introduced punishment threat hinders, norm compliance. Here, we used a dictator game and fMRI to investigate how intention modulates the effect of punishment threat on norm compliance and the substrates of this modulation. We also investigated whether this modulation can be influenced by brain stimulation. Human participants divided an amount of money between themselves and a partner. The partner (intentionally) or a computer program (unintentionally) decided to retain or waive the right to punish the participant upon selfish distribution. Compared with the unintentional condition, participants allocated more when the partner intentionally waived the power of punishment, but less when the partner retained power. The right lateral orbitofrontal cortex (rLOFC) showed higher activation when the partner waived compared with when the computer waived or when the partner retained the power. The functional connectivity between the rLOFC and the brain network associated with intention/mentalizing processing was predictive of the allocation difference induced by intention. Moreover, inhibition or activation of the rLOFC by brain stimulation decreased or increased, respectively, the participants' reliance on the partner's decision during monetary allocation. These findings demonstrate that the perceived intention of punishment threat plays a crucial role in norm compliance and that the LOFC is causally involved in the implementation of intention-based cooperative decisions.

Key words: intention; lateral orbitofrontal cortex; norm compliance; punishment threat; tDCS

Introduction

Social norms are widely shared rules about what constitutes appropriate behavior in social interactions (Bicchieri, 2006). Pun-

ishment is a ubiquitously adopted approach in human society to enforce norm compliance beyond the recipients' voluntary action. Recent studies, however, provide divergent evidence concerning the effect of punishment threat on norm compliance. Studies reveal that participants achieve a higher level of norm compliance when punishment threat is present than when it is absent (Ehr and Gächter, 2005; Spitzer et al., 2007).

evidence also shows that punishment threat under certain circumstances hinders norm compliance. For example, in the trust game, the trustee returns less money to the investor when the investor imposes a punishment threat on the trustee (Furstedt, Rockenbach, 2003; Gneezy and Rustichini, 2000; Oudiz et al., 2009; Li et al., 2009). The neural activity also shows contrasting patterns (Spitzer et al., 2007) and that activations in the lateral orbitofrontal cortex (LOFC) and dlPFC were positively correlated with individuals' increase in norm compliance when a punishment threat was present. In contrast, (Oudiz et al., 2009) observed decreased activations in the LOFC and ventromedial prefrontal cortex (vmPFC) when punishment threat was present.

Closer examination of previous studies reveals that those reporting a detrimental effect typically adopted intentional punishment threat imposed by the interacting partner on behalf of his/her own interest (Fehr and Rockenbach, 2003; Oudiz et al., 2009) whereas those reporting a facilitatory effect involved unintentional punishment threat, which was introduced by an impartial third-party (e.g., computer program) for the sake of fairness (Spitzer et al., 2007; Ruff et al., 2013). However, to our knowledge, no studies have investigated directly the role of intention behind punishment threat in norm enforcement. We hypothesized that the seemingly contradicting findings concerning the role of punishment threat could be reconciled if we take into account the intention behind the threat (Radey et al., 2012; Koster-Hale et al., 2013).

Of particular interest is the orbitofrontal cortex, a structure consistently implicated in computing social value and guiding social decision making (Rushworth et al., 2003; Rudebeck and Murray, 2008). We hypothesized that the LOFC may synthesize information about the presence of punishment threat and the intention by which it is imposed or forgone to form a unified signal that guides compliance behavior (Carron-Bell-Meiklejohn et al., 2012).

To test our hypotheses, we manipulated the presence of punishment threat (Waive vs Retain) and the intention behind the threat (Intentional vs Unintentional) in a modified dictator game. By conducting an fMRI and two high-definition transcranial direct current stimulation (HD-tDCS) experiments, we examined the modulation of the neural processes of punishment threat by the intention behind such a threat. We were specifically interested in the role of the LOFC in mediating the influence of the perceived intention on norm compliance because this structure showed opposite effects when the threat was unintentional (Spitzer et al., 2007) or intentional (Li et al., 2009).

Materials and Methods

Participants

fMRI experiment. Thirty-five graduate and undergraduate students participated in the fMRI scanning. Ten were excluded (1 of them was transferred 0 yuan to the partner; 7 of them did not believe that they interacted with different human partners, as indicated in the postscan manipulation check; 2 of them had excessive head movements in rotation or >3 mm in translation during the scanning), leaving 25 participants for data analysis (age range: 18–27 years, mean age: 23.2 years; 14 female). Due to technical problems, postscan questionnaires and data were available for only 19 of these participants. We tested the robustness of online behavioral measures and postscan questionnaires (e.g., emotion ratings) in an independent sample of participants (see below).

Behavioral validation experiment. To test the stability of the behavioral patterns that we observed in the fMRI experiment, we performed a behavioral experiment with the same procedure as the fMRI experiment

in an independent sample of 24 participants (age range: 18–24 years, mean age: 19.9 years; 9 female).

Brain stimulation experiments. Forty-three graduate and undergraduate students participated in the tDCS experiments. One group of these participants ($n = 22$, age range: 19–25 years, mean age: 21.2; 16 female) received cathodal and sham treatment in 2 experimental sessions separated by 42 d, whereas the other group received anodal and sham treatment, also in 2 experimental sessions. One participant of the latter group failed to show up for the second session, leaving 20 participants in the anodal experiment (age range: 18–25 years, mean age: 21.0 years; 10 female).

None of the participants reported any history of psychiatric, neurological, or cognitive disorders. Informed written consent was obtained from each participant before the experiments. The study was performed in accordance with the Declaration of Helsinki and was approved by the Ethics Committee of the Department of Psychology, Peking University.

Design and procedures

The experiment was a 2 (decider: Computer vs Partner) by 2 (threat: Waive vs Retain) within-participant factorial design. A modified repeated one-shot dictator game was used, in which the participant allocated 20 yuan (\$3.50) between him/herself and a randomly paired partner (chosen from three confederates). In each round, the computer (in the unintentional conditions) or the paired partner (in the intentional conditions) decided to retain or to waive the punishment threat (4 yuan) before the participant made the allocation. In addition, the participants were told that, in each round, the paired partner decided a minimal amount of allocation that he/she would like to accept, although this amount would not be communicated to the participant. If the amount allocated to the partner was less than the minimum and the punishment threat was retained (either by the partner or by the computer), then the punishment would be executed and 4 yuan would be subtracted from the participants' payoff for the current trial. We did not provide trial-by-trial feedback concerning payoff to the participants to prevent the participants from learning a specific behavioral strategy. The amount allocated to the paired player was a measure of the participant's norm compliance. Upon arrival at the laboratory, the participant was introduced to three same-sex strangers, who were in fact confederates of the experiment. The participant was assigned the role of allocator and the confederates were always the responders. The participant was made to believe that each trial he/she would play the game through internet with a randomly paired responder in another room. The participant was told that, after the experiment, one of his/her decisions would be chosen randomly and actualized. We also told the participant that, because no one knew which trial would be selected in the end, the best strategy for him/her was to treat each trial equally seriously.

Each trial began with the presentation of a white fixation against a black background lasting for 4000–6000 ms. Then, a cue of the total allocation amount (a picture of a 20 yuan bill) was presented for 2000 ms, followed by a sentence indicating that the decider (partner or computer) was considering whether to retain punishment threat. This sentence remained on the screen for 2000–5000 ms. Then, the decision (to retain or to waive), together with a cue of the decider (a picture of either a computer or a human silhouette), was presented on the screen for 3000 ms. Finally, after a 2000–4000 ms fixation, a distribution screen was presented and the participant was asked to make the allocation to himself or to the partner (with a step of 2 yuan) by pressing 2 buttons to increase or decrease the amount to be allocated to the partner (with a step of 2 yuan) before pressing another button to confirm the allocation. Button mapping was counterbalanced across participants. The initial amount on the side of the participant was either 0 or 20 yuan and was balanced within each condition. The participant had up to 10 s for the allocation.

The experiment consisted of two blocks of 32 trials each. In each of the four experimental conditions contained 16 trials. Unknown to the participant, the sequence of trials was predetermined by a computer program. The 32 trials in the first block were pseudorandomized with the restriction that no more than three consecutive trials were from the same condition and the second block used the inverted se-

corresponding to the contrast Partner_Retain – Computer_Retain (i.e., intentional punishment threat hinders norm compliance) and Partner_Waive Computer_Waive (i.e., refraining from the threat of punishment facilitates norm compliance). To test the possibility that the strength of such functional connectivity is modulated by individuals' susceptibility to the intention effect, we added the difference in allocation corresponding to each of these contrasts as a group-level covariate. We then used the one-sample *t*-test in SPM8 to perform statistical analysis. The statistic threshold was the same as indicated above.

Brain stimulation experiment

To test the causal role of the rLOFC in mediating the influence of intention on punishment threat, we performed two brain stimulation experiments using HD-tDCS. The first group of participants ($n = 22$) received cathodal stimulation and sham stimulation in two experiment sessions. Half of the participants received cathodal stimulation over the rLOFC in the first experiment day and received sham stimulation over the same area in the second experiment day. The other half of the participants received the reversed stimulation protocol. The second group of participants ($n = 20$) received anodal stimulation and sham stimulation in two experiment sessions. Similar to the cathodal experiment, half of these participants received anodal stimulation over the rLOFC in the first experiment day and received sham stimulation over the same area in the second experiment day. The other half of the participants received the reversed stimulation protocol. Therefore, both of the two HD-tDCS experiments used a within-participant design; moreover, to avoid carry-over effects of brain stimulation, sessions were separated by at least 24 h for each participant. The behavioral protocol was identical to the fMRI experiment.

HD stimulation was delivered using a multichannel stimulation adapter (Soterix Medical, Model C3) connected to the constant current stimulator (Soterix Medical, Model 1300-A). An montage consisting of five sintered Ag/AgCl ring electrodes was used and these electrodes were arranged on the skull in a ring configuration as suggested by the previous literature (Munoz et al., 2011). The electrodes were held in place in plastic electrode holders in a fitted cap (EASYCAP). The electrode holders were filled with SignaGel, creating a gel contact of $\sim 4 \text{ cm}^2$ per electrode. The position of the electrode was identified and adjusted using HD-Explore software (Soterix Medical), which uses a finite-element-method modeling approach to quantify electric field intensity throughout the brain (Datta et al., 2009; Mochowski et al., 2011; Kempe et al., 2014). The locations of the electrodes were chosen by selecting the 10–20 EEG sites that would optimally target the rLOFC in our fMRI study. Therefore, we selected central electrode as FP2 in the 10–20 EEG coordinate system and surrounded it with three return electrodes at F2, F8, Fp1, and one return electrode at the lower eyelid (each at a distance of 6 cm from the central electrode). For active anodal/cathodal stimulation, participants received a constant current of 2.0 mA for min. Stimulation started 8 min before the task and was delivered during the entire course of the task (20 min), with an additional 30 s ramp-up at the beginning of stimulation and 30 s ramp-down at the end. For the sham stimulation, the initial 30 s ramp-up was immediately followed by the 30 s ramp-down and there was no stimulation for the rest of the session. For both the experimental and sham stimulation conditions, participants felt a little uncomfortable initially, but were unaware of stimulation before the task started.

the behavioral validation experiment ($F_{(1,23)} = 10.83, p < 0.001$). Retain > Waive revealed activations in the dmPFC, thalamus, Pairwise comparison showed that, compared with the Computer_Waive condition, participants allocated significantly more to the partner in the Partner_Waive condition ($F_{(1,23)} = 4.85, p < 0.05$); compared with the Computer_Retain condition, participants allocated less to the partner in the Partner_Retain condition ($F_{(1,23)} = 3.33, p = 0.081$).

For the emotional ratings (Fig. 2D), we averaged the ratings of happiness, benevolence, and gratitude to form an indicator of positive affect and the ratings of sadness, anger, fear, and hostility to form an indicator of negative affect. We then performed a repeated-measures ANOVA with emotional valence (Positive vs Negative), Decider (Partner vs Computer), and Threat (Retain vs Waive) as within-participant factors. No main effects were observed. However, a significant interaction was observed between Threat and Decider ($F_{(1,18)} = 28.94, p < 0.001$). Pairwise comparison showed that the positive affect was higher in the Partner_Waive condition than in the Computer_Waive and the Partner_Retain conditions ($F > 37, p < 0.001$). For the negative affect, the interaction was significant ($F_{(1,18)} = 7.12, p < 0.05$). The negative affect was higher in the Partner_Retain condition than in the Computer_Retain and the Partner_Waive conditions ($F > 5, p < 0.05$). Moreover, we performed a two-way ANOVA of the ratings of perceived trust. The interaction was significant ($F_{(1,18)} = 33.52, p < 0.001$). Pairwise comparison showed that the perceived trust was higher in the Partner_Waive condition than in the Computer_Waive condition ($F_{(1,18)} = 68.16, p < 0.001$) and the Partner_Retain condition ($F_{(1,18)} = 32.03, p < 0.001$).

Again, the postexperiment ratings of behavioral validation experiment replicated the behavioral data of the fMRI experiment. For positive emotions, the Decider-by-Threat interaction was significant ($F_{(1,23)} = 49.79, p < 0.001$). Pairwise comparison showed that positive affect was higher in the Partner_Waive condition than in the Computer_Waive and the Partner_Retain conditions ($F > 73, p < 0.001$). For the negative affect, the two-way interaction was marginally significant ($F_{(1,23)} = 3.80, p = 0.064$). The negative affect was higher in the Partner_Retain condition than in the Computer_Retain and the Partner_Waive conditions ($F > 11, p < 0.01$). For perceived trust, the Decider-by-Threat interaction was significant ($F_{(1,23)} = 22.70, p < 0.001$). The perceived trust was higher in the Partner_Waive condition than in the Computer_Waive condition ($F_{(1,23)} = 52.18, p < 0.001$) and the Partner_Retain condition ($F_{(1,23)} > 27.14, p < 0.001$). Together, these results strongly indicate that intentionally introducing punishment threat elicits strong negative emotions, whereas intentionally waiving punishment threat elicits strong positive emotions such as gratitude and the feeling of being trusted.

Whole-brain analysis of the neuroimaging data

When the decision was to retain the punishment threat, the participants were facing certain danger and provocation regardless of whether it was made by the partner or by the computer program. Previous studies have shown that several brain areas related to mentalizing (e.g., dmPFC, TPJ) and affective salience (e.g., thalamus, insula, caudate) are recruited in situations of active aggression and hostility (Kiehl et al., 2001; Beyer et al., 2015). Consistent with these findings, the main effect contrast Retain > Waive revealed activations in the dmPFC, thalamus, insula, caudate, and TPJ (Fig. 3A). To test our hypothesis concerning the modulation of intention on the effect of punishment threat, we examined the interaction contrast (Partner_Waive > Computer_Waive) > (Partner_Retain > Computer_Retain). This contrast revealed activations in the bilateral LOFC (left LOFC: MNI coordinates $[-42, 32, 1]$, cluster size $7, t_{(24)} = 3.66$; rLOFC: MNI coordinates $[42, 35, -5]$, cluster size $72, t_{(24)} = 3.85$; Fig. 3B). Given that we did not observe an interaction in the vmPFC at the current threshold level, we performed an ROI-based analysis within a predefined vmPFC ROI (small volume correction within 8 mm-radius sphere around $[4, -4, -4]$, the coordinates reported in Li et al., 2009). This analysis did reveal a significantly activated cluster (MNI coordinates $[56, -8]$; cluster size 25 voxels; $t_{(24)} = 3.32$; peak-level $p_{FWE} < 0.05$; Fig. 3C). The reversed contrast did not reveal any significant clusters. To illustrate the interaction more clearly, we decomposed the interaction into two separate contrasts: Computer_Retain > Computer_Waive, which corresponded to unintentional punishment threat (Spitzer et al., 2007) and Partner_Waive > Partner_Retain, which corresponded to intentionally withdrawing the punishment threat (Li et al., 2009). The former contrast (Fig. 3C) revealed activation clusters in the left LOFC (MNI coordinates $[-39, 32, 1]$, cluster size $3, t_{(24)} = 4.18$) and the left caudate (MNI coordinates $[-9, 8, 1]$, cluster size 106 , $t_{(24)} = 3.70$). The latter contrast (Fig. 3D) revealed only one activated cluster in the rLOFC (MNI coordinates $[39, 35, -5]$, cluster size $48, t_{(24)} = 3.88$).

ROI-based analysis of the neuroimaging data

To buttress the findings derived from the whole-brain analysis, we performed further analyses for predefined ROIs: the vmPFC and the LOFC. We hypothesized that, if vmPFC activation reflected positive social value (eg, mutual trust) perceived in the dyadic interaction, then it should show higher activation when the partner intentionally waived the punishment threat, an action that may convey trust (Fig. 2B), than when the partner retained the threat. To test this hypothesis, we performed a small volume correction within the vmPFC ROI (8 mm-radius sphere around $[4, -4, -4]$, coordinates reported in Li et al., 2009). This analysis revealed a significantly activated cluster in the vmPFC ROI (MNI coordinates $[3, 56, -8]$; cluster size 17 ; $t_{(24)} = 3.41$; peak-level $p_{FWE} = 0.013$; Fig. 3D). Concerning the rLOFC, we hypothesized that its responses to punishment threat should be modulated by the intentionality behind the threat. Specifically, the rLOFC activation should be higher in the Computer_Retain condition than in the Computer_Waive condition, whereas the opposite pattern should be observed for the Partner conditions. To this end, we performed a small volume correction within the rLOFC ROI (8-mm-radius sphere around $[44, -4]$, coordinates reported in Spitzer et al., 2007). Within this rLOFC ROI, the contrast Computer_Retain > Computer_Waive revealed a significantly activated cluster centered around the MNI coordinates $[51, 38, -2]$ (cluster size 2 ; $t_{(24)} = 2.91$; peak-level $p_{FWE} < 0.05$), while the contrast Partner_Waive > Partner_Retain revealed a significantly activated cluster centered around the MNI coordinates $[39, 35]$ (cluster size 15 ; $t_{(24)} = 3.54$; peak-level $p_{FWE} < 0.01$). Such dissociation confirmed our hypothesis concerning the rLOFC. Moreover, the parameter estimates extracted from the predefined rLOFC and vmPFC ROIs (27 voxels around the coordinates $[39, 35]$ and $[3, 56, -8]$, respectively; Spitzer et al., 2007; Li et al., 2009) for rLOFC and

vmPFC, respectively) exhibited a pattern generally consistent with our findings derived from the small volume correction analysis (Fig. 3E, F). We performed repeated-measures ANOVAs on the parameter estimates and report the statistical details in Table 1. The Decider-by-Threat interaction was significant for both the rLOFC and the vmPFC. Specifically, for the vmPFC, the activation was significantly higher in the Partner_Waive condition than in the Partner_Retain condition (i.e., the same as reported in Li et al., 2009) and was also significantly higher than in the Computer_Waive condition, consistent with the social value representation view of vmPFC function (Ruff and Fehr, 2001). For the rLOFC, the parameter estimates appeared to be higher in the Partner_Waive condition than in the Partner_Retain condition and the parameter estimates appeared to be higher in the Computer_Retain condition than in the Computer_Waive condition, although these differences did not reach statistical significance.

Functional connectivity (PPI) analysis

We performed PPI analyses to test whether the functional connectivity between the mentalizing network and the left vmPFC or the rLOFC was modulated by experimental manipulation and whether such connectivity was predictive of participants' norm compliance behavior. The functional connectivity (for the contrast Partner_Waive > Computer_Waive) between the rLOFC and several brain areas in the typical mentalizing network (e.g., dmPFC, TPJ, and precuneus) was positively correlated with the difference in allocation amount between the Partner_Waive and Computer_Waive conditions (Fig. 4 yellow areas; Table 2).

Similarly, the functional connectivity (for the contrast Partner_Retain > Computer_Retain) between the rLOFC and several brain areas in the typical mentalizing network (e.g., dmPFC, TPJ, and precuneus) was positively correlated with the difference in allocation amount between the Partner_Retain and Computer_Retain conditions (Fig. 4 blue areas; Table 2). No significant result was revealed by the PPI analysis with vmPFC.

Brain stimulation (HD-tDCS) results

For each of the tDCS experiments, we performed a repeated-measures ANOVA with Stimulation Type (Cathodal/Anodal vs Sham), Decider (Computer vs Partner), and threat (Retain vs Waive) as within-participant factors. For the cathodal experiment, the three-way interaction was significant ($F_{(2, 18)} = 5.97, p < 0.05$; Fig. 5A). We then performed a two-way ANOVA focusing on the data in which the partner determined the presence or absence of the punishment threat. The interaction between Stimulation Type and Threat was significant ($F_{(1, 9)} = 11.10, p < 0.01$).




Figure 3. ROI analysis of brain activations. A, B, Brain slices showing activation clusters in the caudate (cau), LOFC, and vmPFC. C, D, Brain slices showing activation clusters in the caudate (cau), LOFC, and vmPFC. E, F, Bar charts showing parameter estimates for the caudate (E) and vmPFC (F) for Computer and Partner conditions across Retain and Waive threats. Error bars represent standard error. Significance markers are shown above the bars.

Table 1. ROI analysis of brain activations

ROI	Computer		Partner	
	$F_{(1, 9)}$	p	$F_{(1, 9)}$	p
cau	2.0	0.18	0.0	0.96
LOFC	9.9	0.01	0.0	0.96
vmPFC	0.0	0.96	0.0	0.96

Psycho-physiological interaction allocation difference as covariate




Figure 4.




Table 2. Brain activations revealed by the PPI covariate (uncorrected at voxel level, cluster-level FWE corrected)

Stim	Threat	MNI	MNI			
			x	y	z	
Cathodal	R	8	2	4	6	
	L	5	8	-8	3	
	R	5	9	3	4	
	L	5	9	-3	-4	
	R	5	9	7	-4	
	L	2	2	-5	-8	3
Anodal	R	8	0	3	-6	
	L	4	3	-5	7	1
	L	9	3	-2	4	0
	L	2	8	-2	4	3
	R	5	5	0	-5	-2
	L	3	0	6	-3	6

0.005). Pairwise comparison showed that, relative to the sham stimulation, the cathodal stimulation decreased the participants' norm compliance behavior (i.e., monetary allocation) when the partner's decision was to waive the punishment threat ($F_{(1,21)} = 4.91p < 0.05$) and increased the allocation when the partner's decision was to retain the punishment threat ($F_{(1,21)} = 5.56p < 0.05$). The same analysis was also applied to the Computer conditions, but neither the main effect nor the interaction was significant.

For the anodal experiment, the three-way interaction was significant ($F_{(1,19)} = 6.00p < 0.05$; Fig. 5B). We then performed a two-way ANOVA focusing on the Partner conditions. The interaction between Stimulation Type and Threat was significant ($F_{(1,19)} = 20.68p < 0.001$). Pairwise comparison showed that, relative to the sham stimulation, the anodal stimulation increased the participants' allocation when the partner's decision

was to waive the punishment threat ($F_{(1,19)} = 8.87p < 0.01$) and decreased the allocation when the partner's decision was to retain the punishment threat ($F_{(1,19)} = 13.57p < 0.005$). The same analysis applied to the Computer conditions revealed neither a significant main effect nor a significant interaction.

To better illustrate and examine the effects of brain stimulation (both inhibition and activation) on intentional/unintentional norm enforcement, we calculated the effect of punishment threat (i.e., the amount transferred in the Waive condition minus the amount transferred in the Retain condition) in the intentional (Partner) and unintentional (Computer) contexts for both the cathodal and anodal groups (Fig. 5C). We then performed two repeated-measures ANOVAs with Stimulation Type (Cathodal/Anodal vs sham) and Decider (Computer vs Partner) as within-participant factors. For the cathodal group, the interaction between Stimulation Type and Threat was significant ($F_{(1,19)} = 5.96p < 0.05$).

Relative to the sham stimulation, the cathodal stimulation decreased the effect of punishment threat mainly in the intentional context ($F_{(1,21)} = 11.10p < 0.005$), but not in the unintentional context ($F_{(1,21)} = 3.60p = 0.072$). For the anodal group, the interaction between stimulation type and threat was significant ($F_{(1,19)} = 5.99p < 0.05$). Relative to the sham stimulation, the anodal stimulation increased the effect of punishment threat only in the intentional context ($F_{(1,19)} = 20.68p < 0.001$), not in the unintentional context ($F_{(1,19)} < 1, p > 0.1$).

Two features of this pattern are worth noting. First, inhibition and activation of the rLOFC had opposite effects on the participants' norm compliance behavior (i.e., monetary allocation): whereas activation of this area tended to increase the effect of waiving the punishment threat on norm compliance (cf. filled and empty red diamonds in Fig. 5C), inhibition of this area tended to decrease this effect (cf. filled and empty blue diamonds in Fig. 5C). Second, the brain stimulation took effect mainly in the intentional context (cf. difference between filled-empty pairs of the Partner side with its counterparts on the Computer side in Fig. 5C).

Discussion

Our behavioral results demonstrated that the perceived intention moderates the effect of punishment threat on norm compliance behavior. Specifically, we observed a detrimental effect of punishment threat in the intentional context (i.e., partner as decider), consistent with previous studies (and Rockenbach, 2008; Rezaee and Attichini, 2004; et al., 2009). In the unintentional context (i.e., computer as decider), although we did not observe a main effect of punishment threat, as previous studies did (and Gächter, 2005; Spitzer et al., 2007; Ruff et al., 2011), the disappearance of the detrimental effect suggests that intention does play an important role in the effectiveness of punishment threat. The intention underlying punishment threat may influence a key factor in norm compliance behavior: the perceived legitimacy of authority. When an impartial computer program or a third party decides to retain the power to punish the allocator, it is

conceived that the retention of punishment threat is on behalf of the social norms themselves. This argument is supported by both our study, which revealed no detrimental effects on norm compliance, and previous studies, which revealed facilitatory effects on norm compliance (Spitzer et al., 2007; Buff et al., 2013). In contrast, when the partner (i.e., the second party), whose interest is directly affected by the allocation, decides to retain the power to punish the allocator, the purpose of the punishment threat is dubious. It may be perceived, not as a way to maintain justice, but rather as a way to serve selfish interest or to signal distrust, resulting in reduced norm compliance (Kin-son and Villeval, 2008). This argument is supported by our behavioral results and the emotion self-reports indicating that intentional retention of punishment threat elicits stronger negative feelings and less amount of allocation than unintentional retention or intentional waiving of punishment threat. In addition, intention can function in, not only a negative way, but also a positive way. We found that, compared with both unintentional waiving and intentional retention of punishment threat, participants reported stronger positive feelings (e.g., being trusted, more grateful) and allocated more to the partner when the latter intentionally waived the power to punish the former.

Houser et al. (2008) manipulated intention but did not find any effect of intention on norm compliance. The discrepancy between their findings and ours may come from two sources. First, intention was a within-participant factor in their study, but a between-participant factor in our study. Therefore, participants who experienced both intentional and unintentional contexts may exhibit a strengthened contrast between the two contexts, which amplifies the difference between intentional and unintentional punishment threat on the perceived legitimacy of authority. Second, the partner's demand of the allocation was not revealed in our study, but was revealed in Houser et al. (2008). Because the participants clearly knew their partner's demand in Houser et al. (2008), they could easily calculate all of the outcomes (i.e., outcome when keeping the entire investment and being punished vs outcome when returning what the partner demanded) and select the most profitable strategy. Such a experimental setup may drive participants to utility-driven strategies, crowding out the influence of intention.

The average transfer in our study was between 30% and 40% of the endowed amount, even in the punishment threat conditions. This was relatively low compared with previous studies, which usually reported 40% average transfer (Spitzer et al., 2007) or 40–50% transfer (Buff et al., 2013) under punishment threat. The discrepancy may be due to the intensity of punishment threat. In the current study, the intensity was relatively low (4 yuan; the whole allocation endowment was 20 yuan) compared with the previous studies. The intensity of punishment threat may modulate its effect on norm enforcement (Gneezy and Rustichini, 2004) and, intuitively, when the punishment threat is




Figure 5. Behavioral results and brain activity. (A) Behavioral results showing transfer percentages for Compliance and Partner conditions under Cath and Sham groups. (B) Behavioral results showing transfer percentages for Anodal and Sham groups. (C) Brain activity in the LOFC region for Cath and Anodal groups. Error bars represent standard error. Significance markers: * p < 0.05, ** p < 0.01.

large enough, it will dominate people's consideration about norm compliance behavior. The discrepancy between the studies, however, does not eliminate the validity of the intention effect that we observed at small amounts of punishment threat. As Gneezy and Rustichini (2004) stated, we have no evidence to support the hypothesis that the psychological and behavioral factors that drive the reaction to small fines or rewards disappear completely when higher amounts are offered or charged, thus reducing the explanation of behavior to a choice of the most convenient combination of effort and reward. Of particular interest to us is the LOFC, which has been consistently implicated in norm compliance, but has showed opposite site activation patterns depending on whether punishment threat was introduced intentionally or unintentionally (Spitzer et al., 2007; Li et al., 2009). Some propose that the LOFC functions to encode the punishment threat based on the findings that higher LOFC activation is associated with more norm compliance behavior under (unintentional) punishment threat (Spitzer et al., 2007). Our results indicated that this could not be the whole story because the LOFC also showed higher activation when the participant intentionally waived the punishment threat. An alternative interpretation, which fits better with both the previous and the current findings, is that the LOFC integrates information from various sources (e.g., intention, emotion, material interest, etc.) and outputs a decision as to whether to conform to the social norm (Rolls and Grabenhorst, 2007). When the presence or absence of the punishment threat is determined by a nonintentional computer program, it is possible that the decision to conform is dominated by the consideration of material interests; that is, the rational calculation of gains and losses. This argument is supported by findings in the current study (Spitzer et al., 2007)

that the norm compliance behavior and LOFC activation is higher in the presence of punishment threat. When the presence or absence of punishment threat is determined by the partner, it conveys important social information, such as trust or distrust. In such contexts, the LOFC and the participant's norm compliance are sensitive to the social signal behind the punishment threat. This conjecture was buttressed by our brain stimulation data: inhibition or activation of the rLOFC by tDCS decreased or increased the effect of partner's intention on norm compliance behavior. Note that we do not claim the laterality of LOFC because we do not have *a priori* hypothesis. We focused our analysis on the right rather than the left LOFC because of the discrepancy between Spritzer et al. (2009) and Li et al. (2009) on the rLOFC. As can be seen from Figure 3B-D, although both the left and right LOFC were revealed in the interaction contrast, only the rLOFC was activated in both simple effect contrasts: Computer_Retain - Computer_Waive and Partner_Waive - Partner_Retain.

The brain stimulation took effect mainly in the intentional context, not in the unintentional context, suggesting that inhibition or activation of the rLOFC may not affect its function in punishment threat processing, but may disrupt or facilitate its function in interacting with other brain regions that could provide social information (e.g., intention, emotion). This argument was supported by our results showing that the functional connectivity between the rLOFC and the brain network typically associated with intention/mentalizing processing (including dmPFC, TPJ, and precuneus; Ulenberghs et al., 2011) was predictive of the effect of intention on norm compliance. Moreover, the functional connectivity (Partner_Waive - Computer_Waive) between the bilateral insula and the rLOFC positively correlated with the increase in norm compliance behavior. The bilateral insula was found to be associated with the aversion of anticipated guilt by not honoring others (Ulenberghs et al., 2011) which may drive individuals to conform to social norms and to show mutual respect in social interactions (Clonnes and Dufwenberg, 2006). Therefore, it is conceivable that the insula encodes the potential guilt that could arise if the participant fails to honor the partner's trust and benevolence (e.g., in the Partner_Waive condition). Such emotional information may be projected to the LOFC to bias the participants' norm compliance behavior.

Finally, we also found higher activation in the vmPFC when the partner waived the power to punish the participant compared with when the partner retained or when the computer waived such power. This is consistent with Li et al. (2009) in which the vmPFC showed higher activation when the partner voluntarily waived the power to punish the participants. Ample evidence has implicated the vmPFC in computing both social and nonsocial reward values (Ruff and Knutson, 2010; Bartra et al., 2008; Ruff et al., 2011). For example, the act of saving money is valued differently and elicits differential activation in the vmPFC according to whether the saving is for charitable donation (higher social value) or for self-interest (lower social value) (Coper et al., 2010; Hare et al., 2010). We argue that the partner's voluntary waiving of the power to punish (i.e., trust and benevolence) is perceived to be most valuable to the individuals.

In conclusion, by combining an interactive game, fMRI, and HD-tDCS, we demonstrate that intention plays an important role in the effectiveness of punishment threat on norm compliance and that the LOFC is causally involved in the implementation of intention-based cooperative decisions.

References

- Eschenko, McGuire JT, Kable JW (2013) The valuation system: a coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *Neuroimage* 76:412-427. [CrossRef Medline](#)
- Beyer F, Munte TF, Guttlich M, Kramer UM (2015) Orbitofrontal cortex reactivity to angry facial expression in a social interaction correlates with aggressive behavior. *Cereb Cortex* 25:3057-3063. [CrossRef Medline](#)
- Bicchieri C (2006) *The grammar of society: the nature and dynamics of social norms*. New York: Cambridge University.
- Plancher JJ, Bikson M, Frohman H, Reeves ST, Datta A, Bansal V, Madan A, Barth K, George MS (2012) A pilot study of the tolerability and effects of high-definition transcranial direct current stimulation (HD-tDCS) on pain perception. *J Pain* 13:112-120. [CrossRef Medline](#)
- Campbell-Meiklejohn DK, Kanai R, Bahrami B, Bach DR, Dolan RJ, Roepstorff A, Frith CD (2012) Structure of orbitofrontal cortex predicts social influence. *Curr Biol* 22:R123-R124. [CrossRef Medline](#)
- Capassi L, Daquer EM, Zimmermann TJ, Mooshagian E, Parra LC, Rice JK, Barba A, Bikson M, Wassermann EM (2012) A pilot study on effects of 4x1 high-definition tDCS on motor cortex excitability. *Conf Proc IEEE Eng Med Biol Soc* 2012:735-738. [CrossRef Medline](#)
- Carlsmith KM, Darley JM, Robinson PH (2002) Why do we punish?: Deterrence and just deserts as motives for punishment. *J Pers Soc Psychol* 83:984-999. [CrossRef Medline](#)
- Chiong J, Smith A, Dufwenberg M, Sanfey AG (2011) Triangulating the neural, psychological, and economic bases of guilt aversion. *Neuron* 70:660-672. [CrossRef Medline](#)
- Cheung G, Dufwenberg M (2006) Promises and partnership. *Econometrica* 74:1579-1601. [CrossRef](#)
- Cooper JC, Kreps TA, Wiebe T, Pirkil T, Knutson B (2010) When giving is good: ventromedial prefrontal cortex activation for others' intentions. *J Neurosci* 30:1511-1521. [CrossRef Medline](#)
- Darley JM (2009) Morality in the law: the psychological foundations of citizens' desires to punish transgressions. *Annual Review of Law and Social Science* 5:1-23. [CrossRef](#)
- Datta A, Bansal V, Diaz J, Patel J, Reato D, Bikson M (2009) Gyri-precise head model of transcranial direct current stimulation: improved spatial focality using a ring electrode versus conventional rectangular pad. *Brain Stimul* 2:201-207. [CrossRef Medline](#)
- Dickinson D, Villeval M (2008) Does monitoring decrease work effort? The complementarity between agency and crowding-out theories. *Games and Economic Behavior* 63:56-76. [CrossRef](#)
- Jamochowski JP, Datta A, Bikson M, Su Y, Parra LC (2011) Optimized multi-electrode stimulation increases focality and intensity at target. *J Neural Eng* 8:046001. [CrossRef Medline](#)
- Fehr E, Gächter S (2002) Altruistic punishment in humans. *Nature* 415:137-140. [CrossRef Medline](#)
- Fehr E, Rockenbach B (2003) Detrimental effects of sanctions on human altruism. *Nature* 422:137-140. [CrossRef Medline](#)
- Friston KJ, Buechel C, Fink GR, Morris J, Rolls E, Dolan RJ (1997) Psychophysiological and modulatory interactions in neuroimaging. *Neuroimage* 6:218-229. [CrossRef Medline](#)
- Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg MD, Turner R (1998) Event-related fMRI: characterizing differential responses. *Neuroimage* 10:30-40. [CrossRef Medline](#)
- Loewy U, Rustichini A (2004) Incentives, punishment and behavior. In: *Advances in behavioral economics* (Camerer C, Loewenstein G, Rabin M, eds), pp 572-589. Princeton, NJ: Princeton University.
- Haber SN, Knutson B (2010) The reward circuit: linking primate anatomy and human imaging. *Neuropsychopharmacology* 35:463-479. [CrossRef Medline](#)
- Hare TA, Camerer CF, Knopfle DT, Rangel A (2010) Value computations in ventral medial prefrontal cortex during charitable decision making incorporate input from regions involved in social cognition. *J Neurosci* 30:583-590. [CrossRef Medline](#)
- Houser D, Xiao E, McCabe K, Smith V (2008) When punishment fails: research on sanctions, intentions and pardons. *Games* 8:001-016. [CrossRef Medline](#)

ments from neural representations of intentions. *Proc Natl Acad Sci U S A* 110:5648-5653. [CrossRef Medline](#)

Rada S, Guroy B, de Bruijn ER (2012) There's something about a fair split: intentionality moderates context-based fairness considerations in social decision making. *PLoS One* 7:e31493. [CrossRef Medline](#)

Kramer UM, Jansma H, Tempelmann C, Miltner TF (2007) Tit-for-tat: the neural basis of reactive aggression. *Neuroimage* 38:203-211. [CrossRef Medline](#)

Rolls ET, Grabenhorst F (2008) The orbitofrontal cortex and beyond: from affect to decision-making. *Prog Neurobiol* 86:216-244. [CrossRef Medline](#)

Kramer UM, Riba J, Richter S, Miltner TF (2011) An fMRI study on the role of serotonin in reactive aggression. *PLoS One* 6:e27538. [CrossRef Medline](#)

Rudebeck PH, Murray EA (2014) The orbitofrontal oracle: cortical mechanisms for the prediction and evaluation of specific behavioral outcomes. *Neuron* 84:1143-1156. [CrossRef Medline](#)

Kuo HI, Bikson M, Datta A, Minhas P, Paulus W, Kuo MF, Nitsche MA (2013) Comparing cortical plasticity induced by conventional and high-definition 1 ring tDCS: a neurophysiological study. *Brain Stimul* 6:644-648. [CrossRef Medline](#)

Ruff CC, Fehr E (2014) The neurobiology of rewards and values in social decision making. *Nat Rev Neurosci* 15:549-562. [CrossRef Medline](#)

Ruff CC, Ugazio G, Fehr E (2013) Changing social norm compliance with noninvasive brain stimulation. *Science* 342:482-484. [CrossRef Medline](#)

Li J, Xiao E, Houser D, Montague PR (2009) Neural responses to sanction threats in two-party economic exchange. *Proc Natl Acad Sci U S A* 106:16835-16840. [CrossRef Medline](#)

Rushworth MF, Noonan MP, Boorman ED, Walton ME, Behrens TE (2011) Frontal cortex and reward-guided learning and decision-making. *Neuron* 70:1054-1069. [CrossRef Medline](#)

Minhas P, Bansal V, Patel J, Ho JS, Diaz J, Datta A, Bikson M (2010) Electrodes for high-definition transcutaneous DC stimulation for applications in drug delivery and electrotherapy, including tDCS. *J Neurosci Methods* 190:188-197. [CrossRef Medline](#)

Spring XW, Dong ZY, Long XY, Li SF, Zuo XN, Zhu CZ, He Y, Yan CG, Zang YF (2011) REST: A toolkit for resting-state functional magnetic resonance imaging data processing. *PLoS One* 6:e25931. [CrossRef Medline](#)

Molenberghs P, Johnson H, Henry JD, Mattingley JB (2016) Understanding the minds of others: A neuroimaging meta-analysis. *Neurosci Biobehav Rev* 65:276-291. [CrossRef Medline](#)

Spitzer M, Fischbacher U, Herrnberger B, Groll F, Fehr E (2007) The neural signature of social norm compliance. *Neuron* 56:185-196. [CrossRef Medline](#)